

Conformal compactification and cycle-preserving symmetries of spacetimes

Francisco J. Herranz[†] and Mariano Santander[‡]

[†] *Departamento de Física, Escuela Politécnica Superior
Universidad de Burgos, E-09006 Burgos, Spain*

[‡] *Departamento de Física Teórica, Facultad de Ciencias
Universidad de Valladolid, E-47011 Valladolid, Spain*

Abstract

The cycle-preserving symmetries for the nine two-dimensional real spaces of constant curvature are collectively obtained within a Cayley–Klein framework. This approach affords a unified and global study of the conformal structure of the three classical Riemannian spaces as well as of the six relativistic and non-relativistic spacetimes (Minkowskian, de Sitter, anti-de Sitter, both Newton–Hooke and Galilean), and gives rise to general expressions holding simultaneously for all of them. Their metric structure and *cycles* (lines with constant geodesic curvature that include geodesics and circles) are explicitly characterized. The corresponding cyclic (Möbius-like) Lie groups together with the differential realizations of their algebras are then deduced; this derivation is new and much simpler than the usual ones and applies to any homogeneous space in the Cayley–Klein family, whether flat or curved and with any signature. Laplace and wave-type differential equations with conformal algebra symmetry are constructed. Furthermore, the conformal groups are realized as matrix groups acting as globally defined linear transformations in a four-dimensional ‘conformal ambient space’, which in turn leads to an explicit description of the ‘conformal completion’ or compactification of the nine spaces.

1 Introduction

The role of conformal groups in physics can hardly be overestimated. Taking aside conformal invariance in quantum field theory, this role appears even at a rather basic level. In special relativity, spacetime is a flat (zero curvature) pseudo-Riemannian space $ISO(3,1)/SO(3,1)$ whose kinematical motion group is the Poincaré group $ISO(3,1)$; its ‘angle’-preserving transformations generate a group, the (Minkowskian) conformal group $SO(4,2)$, which has dimension 15 and contains as a subgroup the Poincaré group. This conformal group is the maximal invariance group of the vacuum Maxwell equations [1, 2, 3, 4], and in general, of a large number of equations in different areas, such as, for instance, all equations describing zero-mass systems [5, 6, 7, 8, 9].

The group of conformal transformations of the N -dimensional (ND) Euclidean space $\mathbf{E}^N$ was firstly found by Liouville (for an explicit construction see [10]) and is also well known. This group can be obtained from two equivalent approaches. The ‘conformal’ method is to obtain transformations $\phi : \mathbf{E}^N \rightarrow \mathbf{E}^N$ which carry the Euclidean metric g into another proportional to it: $g' \equiv \phi^*g = \mu(x)g$, where $\mu(x)$ is a function of the coordinates; these preserve the angle between any two crossing curves, and generate the Euclidean conformal group. The ‘hypersphere’ method is to look for (local) transformations $\mathbf{E}^N \rightarrow \mathbf{E}^N$ which carry hyperspheres into hyperspheres, including hyperplanes as limit cases; this approach naturally requires to complete $\mathbf{E}^N$ in order to have these transformations globally defined, and has the advantage of showing clearly the structure of the ‘hyperspherical’ group of $\mathbf{E}^N$ [11]; it was introduced by Lie and Darboux. Both constructions lead to identical results except for the 2D case, where the conformal angle-preserving group is infinite-dimensional, while the circle-preserving one (so called circular transformations) has dimension 6. In the generic ND case, the transformations obtained through either method generate, as is well known, a finite-dimensional Lie group isomorphic to $SO_0(N+1,1)$, that is, the connected component of the identity of $O(N+1,1)$ which appears when discrete reflections and inversions are also considered. The corresponding Lie algebra is spanned by the usual Euclidean generators of translations and rotations together with some new generators of dilations and specific conformal transformations.

The Minkowskian and Euclidean spaces are two important particular instances within the family of *flat* symmetric homogeneous spaces $\mathbf{R}^{p,q} \equiv ISO(p,q)/SO(p,q)$. The conformal group of $\mathbf{R}^{p,q}$ is isomorphic to $SO_0(p+1, q+1)$ (inversions not included) [10, 12]; its transformations preserve the ‘angle’ between any two curves and geometrically they carry $\mathbf{R}^{p,q}$ -hyperspheres into themselves. In this context it is rather natural to inquire which are the conformal groups of the homogeneous Riemannian and pseudo-Riemannian spaces with *non-zero* constant curvature (for $N = 2$ they are the sphere $\mathbf{S}^2$, the hyperbolic plane $\mathbf{H}^2$ and the two de Sitter ‘spheres’ $\mathbf{AdS}^{1+1}$ and $\mathbf{dS}^{1+1}$), and also to analyze the conformal groups of the (contracted) cases with a degenerate metric (for $N = 2$, these are the $(1+1)$ D Galilean $\mathbf{G}^{1+1}$ and the two Newton–Hooke $\mathbf{NH}_\pm^{1+1}$ spacetimes). The above spaces together with the Euclidean $\mathbf{E}^2$ and the Minkowskian $\mathbf{M}^{1+1}$ spaces constitute the nine quasi-orthogonal homogeneous spaces or Cayley–Klein (CK) spaces [13, 14, 15, 16] in two dimensions.

The CK groups share many properties, even if the CK family comprises simultaneously simple as well as non-simple (but still quasi-simple) orthogonal Lie groups [17, 18, 19]. Further to the groups and algebras themselves, this CK framework allows a unified and global approach to the study of the associated homogeneous spaces (including the

relativistic and non-relativistic spacetimes of constant curvature). General expressions parametrized through two real coefficients κ_1, κ_2 and holding simultaneously for all nine 2D CK groups/spaces can be obtained; for each specific case they follow simply by particularizing κ_1, κ_2 . For instance, their corresponding Drinfeld–Jimbo quantum deformations [14, 20], superintegrable systems [21], trigonometry [16], etc. have already been studied by following this ‘universal’ approach which displays clearly some family properties hidden in a case-by-case analysis. The logic behind this approach can be described as ‘extended meaning, simpler proofs’.

The aim of this paper is twofold. First, to study in detail the *cycle-preserving* transformations of the nine 2D CK spaces (the finite dimensional part of the full set of conformal transformations in 2D), paying special attention to the six spacetimes. Second, we propose an explicit description, covering the nine CK spaces in the same run, of their corresponding inversive or conformal compactification, where cycle-preserving transformations can be defined as global transformations. We present a joint view of the structure of these conformal compactifications together with their relationships and the way the initial space is embedded into its compactified. Thus we recover many known results while adopting a fresh perspective we provide a clear view of the behaviour of the compactification when either curvature vanishes or metric degenerates.

In section 2 we describe the nine 2D CK spaces, whose metric structure is studied in section 3 by introducing three sets of geodesic coordinates useful in our further construction. In section 4 we obtain the equations of *cycles* as lines with constant geodesic curvature that include, as particular cases, geodesics, equidistants, horocycles and circles; these complete the results obtained in [16] concerning trigonometry. Next we give a simple and apparently new derivation of cycle-preserving transformations in section 5. In this way, the conformal Lie groups together with the differential realizations of their corresponding conformal algebras are obtained. Differential equations with conformal algebra symmetry are constructed in section 6; these results not only cover the usual 2D Laplace and $(1+1)$ D wave equations, but also encompass their non-zero curvature versions with the associated symmetry operators. The ‘conformal completion’ of the 2D CK spaces is developed in section 7 through a realization of their conformal groups acting as globally defined linear transformations in a 4D ambient space; this extends to all the nine CK spaces the conformal compactification which is already familiar for the Euclidean and Minkowskian spaces but which also makes sense for the curved and degenerate spaces. Some remarks close the paper.

A large amount of information relative to cycle-preserving conformal transformations is made available in completely explicit and tabular form, so that it is easily retrievable. As a byproduct of our construction, a clear and global picture of the conformal symmetries for $(1+1)$ D relativistic and non-relativistic spacetimes with constant curvature is obtained. Along the paper we freely interpret the ‘geometrical’ results by ‘translating’ them into the kinematical language (as in tables 3, 4, 6, 8 and 10).

2 The nine two-dimensional Cayley–Klein spaces

To begin with we briefly recall the algebraic structure of the nine 2D CK spaces (for more details see [16]). Their motion groups are collectively denoted $SO_{\kappa_1, \kappa_2}(3)$ where κ_1, κ_2 are two real coefficients. The commutation relations of the CK algebra $so_{\kappa_1, \kappa_2}(3)$ in the basis

$\{P_1, P_2, J_{12}\}$ and the Casimir invariant read:

$$[J_{12}, P_1] = P_2 \quad [J_{12}, P_2] = -\kappa_2 P_1 \quad [P_1, P_2] = \kappa_1 J_{12} \quad (2.1)$$

$$\mathcal{C} = \kappa_2 P_1^2 + P_2^2 + \kappa_1 J_{12}^2. \quad (2.2)$$

Notice that κ_1, κ_2 can be reduced to $+1, 0, -1$ by rescaling the generators. The *plane* (as the space of points) corresponds to the 2D symmetric homogeneous space

$$S_{[\kappa_1], \kappa_2}^2 = SO_{\kappa_1, \kappa_2}(3)/SO_{\kappa_2}(2) \quad SO_{\kappa_2}(2) = \langle J_{12} \rangle \quad (2.3)$$

hence the generator J_{12} leaves a point O (the origin) invariant, thus acting as the rotation around O , while P_1, P_2 generate translations that move O along two basic directions.

The curvature and metric signature of these CK spaces are determined by κ_1, κ_2 : the space $S_{[\kappa_1], \kappa_2}^2$ has a canonical *metric* which at O comes from the Casimir; as the space is homogeneous the metric is determined once given at any point. This metric turns out to have *constant curvature* κ_1 (written in square brackets in the space notation). At the origin O the metric matrix is $\text{diag}(1, \kappa_2)$ in the basis P_1, P_2 of the tangent space; therefore κ_2 determines the metric *signature*. We display the nine 2D CK spaces in table 1; any vanishing coefficient κ_i can be interpreted as an Inönü–Wigner contraction limit and corresponds to either vanishing curvature ($\kappa_1 \rightarrow 0$) or degenerating metric ($\kappa_2 \rightarrow 0$).

Spacetimes with constant curvature [22] appear in this scheme. If $\{P_1, P_2, J_{12}\}$ are read as generators of time translations, space translations and boosts, respectively, the six CK groups with $\kappa_2 \leq 0$ (second and third rows of table 1) are the (kinematical) motion groups of $(1+1)$ D spacetimes. According to the values of (κ_1, κ_2) we find in table 1:

- Three ‘absolute-time’ spacetimes for $\kappa_2 = 0$: oscillating Newton–Hooke $\mathbf{NH}_+^{1+1} (+, 0)$, Galilean $\mathbf{G}^{1+1} (0, 0)$ and expanding Newton–Hooke $\mathbf{NH}_-^{1+1} (-, 0)$ (we denote $ISO(1) \equiv \mathbb{R}$). These are non-relativistic spacetimes with a degenerate Riemannian metric of signature type $\text{diag}(+, 0)$.
- Three ‘relative-time’ spacetimes for $\kappa_2 < 0$: anti-de Sitter $\mathbf{AdS}^{1+1} (+, -)$, Minkowskian $\mathbf{M}^{1+1} (0, -)$ and de Sitter $\mathbf{dS}^{1+1} (-, -)$, with a Lorentzian (indefinite) metric with signature type $\text{diag}(+, -)$.

In this kinematical context, the coefficients κ_i are related to the universe time radius τ and speed of light c by

$$\kappa_1 = \pm 1/\tau^2 \quad \kappa_2 = -1/c^2. \quad (2.4)$$

The curvature κ_1 may also be considered as a cosmological constant. The contraction $\kappa_1 \rightarrow 0$ corresponds to the flat limit $\tau \rightarrow \infty$; likewise the contraction $\kappa_2 \rightarrow 0$ is the non-relativistic limit $c \rightarrow \infty$. The three remaining spaces with $\kappa_2 > 0$ have no direct kinematical interpretation (unless one passes to imaginary Euclidean time); they are the three classical Riemannian spaces of constant curvature with definite positive metric with signature type $\text{diag}(+, +)$.

3 Metric structure and coordinate systems

3.1 Matrix realization of the Cayley–Klein groups

Let us consider the following 3D real matrix representation of $so_{\kappa_1, \kappa_2}(3)$:

$$P_1 = -\kappa_1 e_{01} + e_{10} \quad P_2 = -\kappa_1 \kappa_2 e_{02} + e_{20} \quad J_{12} = -\kappa_2 e_{12} + e_{21} \quad (3.1)$$

where e_{ij} is a 3D matrix with a single non-zero entry 1 at row i and column j ($i, j = 0, 1, 2$). The exponential of these matrices leads to one-parametric subgroups of $SO_{\kappa_1, \kappa_2}(3)$:

$$\begin{aligned} e^{\alpha P_1} &= \begin{pmatrix} C_{\kappa_1}(\alpha) & -\kappa_1 S_{\kappa_1}(\alpha) & 0 \\ S_{\kappa_1}(\alpha) & C_{\kappa_1}(\alpha) & 0 \\ 0 & 0 & 1 \end{pmatrix} & e^{\gamma J_{12}} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & C_{\kappa_2}(\gamma) & -\kappa_2 S_{\kappa_2}(\gamma) \\ 0 & S_{\kappa_2}(\gamma) & C_{\kappa_2}(\gamma) \end{pmatrix} \\ e^{\beta P_2} &= \begin{pmatrix} C_{\kappa_1 \kappa_2}(\beta) & 0 & -\kappa_1 \kappa_2 S_{\kappa_1 \kappa_2}(\beta) \\ 0 & 1 & 0 \\ S_{\kappa_1 \kappa_2}(\beta) & 0 & C_{\kappa_1 \kappa_2}(\beta) \end{pmatrix} \end{aligned} \quad (3.2)$$

where we have introduced the cosine $C_\kappa(x)$ and sine $S_\kappa(x)$ functions [14, 15, 16, 20]:

$$C_\kappa(x) = \begin{cases} \cos \sqrt{\kappa} x & \kappa > 0 \\ 1 & \kappa = 0 \\ \cosh \sqrt{-\kappa} x & \kappa < 0 \end{cases} \quad S_\kappa(x) = \begin{cases} \frac{1}{\sqrt{\kappa}} \sin \sqrt{\kappa} x & \kappa > 0 \\ x & \kappa = 0 \\ \frac{1}{\sqrt{-\kappa}} \sinh \sqrt{-\kappa} x & \kappa < 0 \end{cases} \quad (3.3)$$

From them, we define the ‘versed sine’ (or versine) $V_\kappa(x)$ and the tangent $T_\kappa(x)$:

$$V_\kappa(x) = \frac{1}{\kappa}(1 - C_\kappa(x)) \quad T_\kappa(x) = \frac{S_\kappa(x)}{C_\kappa(x)}. \quad (3.4)$$

These curvature-dependent functions coincide with the circular and hyperbolic ones for $\kappa = 1$ and $\kappa = -1$, respectively; the contracted case $\kappa = 0$ gives rise to the parabolic or Galilean functions: $C_0(x) = 1$, $S_0(x) = x$ and $V_0(x) = x^2/2$. Identities for the trigonometric functions (3.3)–(3.4) will be necessary in the computations carried out throughout the paper; these relations can be found in the appendix of [16] so that we omit them here. We will also need their derivatives: if we denote their corresponding inverse functions with the prefix ‘arc-’, it can be shown that [15]:

$$\begin{aligned} \frac{d}{dx} C_\kappa(x) &= -\kappa S_\kappa(x) & \frac{d}{dx} \text{arc}C_\kappa(x) &= \frac{-1}{\kappa \sqrt{\frac{1}{\kappa}(1-x^2)}} \\ \frac{d}{dx} S_\kappa(x) &= C_\kappa(x) & \frac{d}{dx} \text{arc}S_\kappa(x) &= \frac{1}{\sqrt{1-\kappa x^2}} \\ \frac{d}{dx} T_\kappa(x) &= \frac{1}{C_\kappa^2(x)} & \frac{d}{dx} \text{arc}T_\kappa(x) &= \frac{1}{1+\kappa x^2} \\ \frac{d}{dx} V_\kappa(x) &= S_\kappa(x) & \frac{d}{dx} \text{arc}V_\kappa(x) &= \frac{1}{\sqrt{2x-\kappa x^2}}. \end{aligned} \quad (3.5)$$

By taking into account the realization (3.2), the CK group $SO_{\kappa_1, \kappa_2}(3)$ can be seen as a group of linear transformations in an ambient space $\mathbb{R}^3 = (x^0, x^1, x^2)$, acting as the group of isometries of a bilinear form Λ with matrix:

$$\Lambda = \text{diag}(1, \kappa_1, \kappa_1 \kappa_2). \quad (3.6)$$

Therefore, a generic group element $X \in SO_{\kappa_1, \kappa_2}(3)$ can be written as a product of the matrices (3.2) satisfying

$$X^T \Lambda X = \Lambda \quad (3.7)$$

where X^T denotes the transpose matrix of X .

The action of $SO_{\kappa_1, \kappa_2}(3)$ on $\mathbb{R}^3$ is linear but not transitive, since it conserves the quadratic form $(x^0)^2 + \kappa_1(x^1)^2 + \kappa_1 \kappa_2(x^2)^2$ provided by Λ . In this action the subgroup

$SO_{\kappa_2}(2) = \langle J_{12} \rangle$ is the isotropy subgroup of the point $O \equiv (1, 0, 0)$ which will be taken as the *origin* in the space $S^2_{[\kappa_1], \kappa_2}$. The action becomes transitive if we restrict to the orbit in $\mathbb{R}^3$ of the point O , which is contained in the ‘sphere’ Σ :

$$\Sigma \equiv (x^0)^2 + \kappa_1(x^1)^2 + \kappa_1\kappa_2(x^2)^2 = 1. \quad (3.8)$$

This orbit is identified with the CK space, and (x^0, x^1, x^2) fulfilling (3.8) are called *Weierstrass coordinates* in $S^2_{[\kappa_1], \kappa_2}$. This scheme includes under a common description all the familiar embeddings (the vector models) of the 2D sphere, hyperbolic plane, anti-de Sitter and de Sitter spacetimes in a linear 3D ambient space with a flat metric of either Euclidean or Lorentzian type. The Weierstrass coordinates allow us to obtain a differential realization of the generators as first-order vector fields in $\mathbb{R}^3$ with $\partial_i = \partial/\partial x^i$ (see (3.1)):

$$P_1 = \kappa_1 x^1 \partial_0 - x^0 \partial_1 \quad P_2 = \kappa_1 \kappa_2 x^2 \partial_0 - x^0 \partial_2 \quad J_{12} = \kappa_2 x^2 \partial_1 - x^1 \partial_2. \quad (3.9)$$

3.2 Metric structure

If both coefficients κ_i are different from zero, $SO_{\kappa_1, \kappa_2}(3)$ is a simple Lie group, and the space $S^2_{[\kappa_1], \kappa_2}$ is naturally endowed with a non-degenerate metric g_0 coming from the non-singular Killing–Cartan form in the Lie algebra $so_{\kappa_1, \kappa_2}(3)$. At the origin, g_0 is given by:

$$g_0(P_1, P_1) = -2\kappa_1 \quad g_0(P_2, P_2) = -2\kappa_1 \kappa_2 \quad g_0(P_1, P_2) = 0. \quad (3.10)$$

When dealing with spaces associated to simple Lie algebras it is customary to define the invariant metric by simply propagating g_0 to the whole space through the group action (after taking out non-essential factors). But we want to cover as well the cases with $\kappa_1 = 0$ where g_0 vanishes identically. To this aim we may take out instead a factor $-2\kappa_1$ out of g_0 in the expression (3.10), and introduce the space *main metric* g_1 from the Killing–Cartan metric g_0 as

$$-2g_1 := g_0/\kappa_1 \quad g_1(P_1, P_1) = 1 \quad g_1(P_2, P_2) = \kappa_2 \quad g_1(P_1, P_2) = 0. \quad (3.11)$$

The metric obtained by propagating g_1 to the whole space is invariant, and is well defined for any value of κ_1 .

The non-generic situation $\kappa_2 = 0$ corresponds to a degenerate metric and is singular. In this case (second row of table 1), the action of $SO_{\kappa_1, 0}(3)$ on $S^2_{[\kappa_1], 0}$ has an invariant foliation, whose set of leaves can be parametrized by $(x^0)^2 + \kappa_1(x^1)^2 = 1 \equiv S^1_{[\kappa_1]}$; each leaf has codimension 1. The restriction of g_1 to each of these leaves vanishes, but $g_2 = \frac{1}{\kappa_2} g_1$ has a non-vanishing and well defined restriction to each leaf, with signature type (+); we will refer to g_2 as the *subsidiary metric*. Summing up, a unified description of the metric structure for the nine 2D CK spaces has the following elements [15, 23]:

- (a) The space $S^2_{[\kappa_1], \kappa_2}$ has a *connection* ∇ (the canonical connection of any symmetric homogeneous space [24]) invariant under $SO_{\kappa_1, \kappa_2}(3)$.
- (b) The space $S^2_{[\kappa_1], \kappa_2}$ has a *hierarchy of two metrics* g_1 and $g_2 = \frac{1}{\kappa_2} g_1$. Generically both metrics are compatible with the connection ∇ . The action of $SO_{\kappa_1, \kappa_2}(3)$ on $S^2_{[\kappa_1], \kappa_2}$ is by isometries of both metrics.
- (c) The main metric g_1 is actually a metric in the true sense (yet it may be degenerate) and has constant curvature κ_1 and signature $\text{diag}(+, \kappa_2)$.

(d) If $\kappa_2 \neq 0$, the connection ∇ is the unique connection compatible with the metric g_1 (thus ∇ is the metric or Levi-Civita connection), and the subsidiary metric g_2 is a true quadratic metric proportional to g_1 .

(e) If $\kappa_2 = 0$ there are more connections compatible with g_1 , but anyhow ∇ is singularized as the canonical one in the homogeneous symmetric space. The subsidiary metric g_2 gives a true metric only in each leaf of the invariant foliation which for $\kappa_2 = 0$ exists in $S^2_{[\kappa_1],0}$ and g_2 has signature (+); in this case the subsidiary metric cannot be overlooked.

In terms of Weierstrass coordinates in the linear ambient space $\mathbb{R}^3$, the two metrics in $S^2_{[\kappa_1],\kappa_2}$ come from the flat ambient metric

$$ds^2 = (dx^0)^2 + \kappa_1(dx^1)^2 + \kappa_1\kappa_2(dx^2)^2 \quad (3.12)$$

in the form

$$(ds^2)_1 = \frac{1}{\kappa_1} ds^2 \quad (ds^2)_2 = \frac{1}{\kappa_2} (ds^2)_1. \quad (3.13)$$

3.3 Geodesic coordinate systems

Now we proceed to introduce three specially relevant coordinate systems of geodesic type in $S^2_{[\kappa_1],\kappa_2}$ (see figure 1). Let us consider the origin $O \equiv (1, 0, 0)$, two (oriented) geodesics l_1, l_2 which are orthogonal through the origin, and a generic point Q with Weierstrass coordinates $\mathbf{x} = (x^0, x^1, x^2)$. By taking into account (3.2), we have:

- If $\mathbf{x} = \exp(aP_1)\exp(yP_2)O$, we will call (a, y) the *geodesic parallel coordinates of type I* of Q . The point Q_1 being the intersection point of the geodesic l'_2 orthogonal to l_1 through Q with l_1 itself, then the first coordinate is the (oriented) distance a between the origin O and the point Q_1 , measured along the geodesic l_1 . The second coordinate is the distance y between Q_1 and Q , measured along the geodesic l'_2 .
- If $\mathbf{x} = \exp(bP_2)\exp(xP_1)O$ we will call (x, b) the *geodesic parallel coordinates of type II* of Q . Now we take as baseline the geodesic l_2 instead of l_1 . Thus, if l'_1 is the geodesic orthogonal to l_2 through Q , and Q_2 is the intersection point of l'_1 and l_2 , then x is the distance between Q_2 and Q measured along l'_1 , while b is the distance between O and Q_2 measured along l_2 .
- The *geodesic polar coordinates* of the point $\mathbf{x} = \exp(\phi J_{12})\exp(rP_1)O$ are (r, ϕ) . The coordinate r is the distance between O and Q measured along the geodesic l joining both points, and ϕ is the angle at O between l and l_1 .

Weierstrass coordinates $\mathbf{x}$ of a generic point Q are displayed in table 2 in the three geodesic coordinate systems; of course the ‘sphere’ condition (3.8) is automatically fulfilled. By introducing these systems in the expressions of the metrics in Weierstrass coordinates, (3.12) and (3.13), we find the main and subsidiary metrics in these intrinsic coordinates. From them we may compute the connexion symbols Γ^i_{jk} . The area element dS in coordinates say u^1, u^2 is $\sqrt{\det g_1/\kappa_2} du^1 \wedge du^2$. Integration of dS over a triangle (easily done in polar coordinates) would lead to the Gauss–Bonnet theorem for the nine CK spaces, recovering the equation (4.29) of [16]. Both metrics, the non-zero Christoffel symbols and the area element are also given in table 2. When the pair (κ_1, κ_2) is particularized to its nine essentially different values, relations appearing in table 2 provide the explicit description of the nine spaces $S^2_{[\kappa_1],\kappa_2}$. This is performed in parallel I and polar coordinates in tables 3 and 4, respectively.




Figure 1: The three geodesic coordinate systems (a, y) , (x, b) and (r, ϕ) of a point Q .

In $\mathbf{E}^2$, geodesic coordinates reduce to the standard cartesian and polar ones (see for instance [25]). In the three flat spaces ($\kappa_1 = 0$), the identities $x = a \equiv x^1$, $y = b \equiv x^2$ hold, so that both parallel I and II systems are the same but this is no longer true for spaces with non-zero curvature ($\kappa_1 \neq 0$).

Each of the systems so obtained covers some domain and may have singularities; both may depend on κ_1, κ_2 . Polar coordinates are always singular at the origin $r = 0$, and in the three relativistic spacetimes with $\kappa_2 = -1/c^2$ they only cover the ‘time-like’ causal cone of O limited by the isotropic lines $x^2 = \pm c x^1$ since $|x^2/x^1| = c |\tanh(\chi/c)| < c$ no matter of the value of the curvature κ_1 . Another ‘type II’ polar coordinate system, with base in the space-like geodesic l_2 instead of the time-like l_1 , might also be introduced; this would cover each half of the exterior of the causal cone in these cases (in the flat case this part is the Rindler space). Similar limitations may happen in parallel I coordinates, which do not cover fully the de Sitter space. When both κ_1, κ_2 are positive, all three coordinate systems cover the whole sphere $\mathbf{S}^2$, and in this case they are essentially equivalent.

The non-relativistic (Newtonian) spacetimes with $c = \infty$ ($\kappa_2 = 0$) have foliations with leaves defined by constant t , and the subsidiary (space) metric is only relevant within them. In parallel I coordinates we find a degenerate main temporal metric $(ds^2)_1 = dt^2$ (providing ‘absolute-time’), and an invariant foliation whose leaves are the ‘absolute-space’ in the moment $t = t_0$ with a non-degenerate subsidiary spatial metric $(ds^2)_2 = dy^2$ defined in each leaf. Notice that the ‘same’ metrics are behind the *three* non-relativistic spacetimes with any curvature; they are distinguished by the canonical connection, which in these cases *cannot* be derived from the metrics.

4 Cycles

In any 2D CK space $S^2_{[\kappa_1], \kappa_2}$, *cycles* are defined as lines with constant geodesic curvature, including *geodesics* as the particular cases with zero geodesic curvature. From a Lie group viewpoint, cycles can also be defined as the orbits of any point by the one-parameter subgroups of the motion group $SO_{\kappa_1, \kappa_2}(3)$. When translated to the vector model in the ambient space $\mathbb{R}^3$, this gives another possible interpretation of geodesics and cycles:

- Geodesics are intersections of the ‘sphere’ Σ (3.8) with planes through the origin in $\mathbb{R}^3$.
- In spaces of *non-zero curvature*, cycles are the intersections of Σ with planes in $\mathbb{R}^3$. The

case of zero curvature, $\kappa_1 = 0$, is special since all plane sections of the ‘sphere’ Σ in $\mathbb{R}^3$ only give geodesics but not cycles.

In this section we characterize cycles and deduce their equations in a direct way. We will assume $\kappa_1 \neq 0$, but the final results will also hold when $\kappa_1 = 0$, as a well defined limit $\kappa_1 \rightarrow 0$ may be meaningfully performed. In the case $\kappa_1 \neq 0$ the equation of a generic cycle is a linear equation in Weierstrass coordinates, where $\alpha, \alpha_0, \alpha_1, \alpha_2$ are constants:

$$\alpha_0 x^0 + \alpha_1 x^1 + \alpha_2 x^2 = \alpha. \quad (4.1)$$

Some properties are better studied in specific coordinate systems in $S^2_{[\kappa_1], \kappa_2}$, and we will freely use the three geodesic coordinates discussed in the previous section. All properties discussed below are coordinate-independent, but in order not to burden the exposition, when using a particular system we will not specify the domains (each coordinate system may not cover all space) nor will we discuss separately their singularities. We start in geodesic parallel I coordinates. By introducing (a, y) of table 2 in (4.1) we obtain the cycle equation in the form:

$$\alpha_0 C_{\kappa_1}(a) + \alpha_1 S_{\kappa_1}(a) = \frac{\alpha}{C_{\kappa_1 \kappa_2}(y)} - \alpha_2 T_{\kappa_1 \kappa_2}(y). \quad (4.2)$$

This equation can be recast in a simpler way that involves the Lambda function $c \rightarrow \Lambda_\kappa(c)$, fully described in appendix A. For any quantity c with label κ , $\Lambda_\kappa(c)$ has the *opposite* label $-\kappa$, and will be denoted simply as $c^\wedge$. If now we consider the Lambda function $\Lambda_{\kappa_1 \kappa_2}(y) \equiv y^\wedge$ of y , whose label is $-\kappa_1 \kappa_2$, the appendix relations (A.10), $1/C_{\kappa_1 \kappa_2}(y) = C_{-\kappa_1 \kappa_2}(y^\wedge)$ and $T_{\kappa_1 \kappa_2}(y) = S_{-\kappa_1 \kappa_2}(y^\wedge)$, allow us to rewrite the cycle equation under the (apparently unknown) form:

$$\alpha_0 C_{\kappa_1}(a) + \alpha_1 S_{\kappa_1}(a) = \alpha C_{-\kappa_1 \kappa_2}(y^\wedge) - \alpha_2 S_{-\kappa_1 \kappa_2}(y^\wedge). \quad (4.3)$$

As is proven in appendix B, the geodesic curvature k_g of the cycle (4.2) (or (4.1)) turns out to be

$$k_g^2 = \kappa_1^2 \alpha^2 \frac{\kappa_2}{\alpha_2^2 + \kappa_2 \alpha_1^2 + \kappa_1 \kappa_2 (\alpha_0^2 - \alpha^2)}. \quad (4.4)$$

The equations of cycles in the two remaining geodesic coordinate systems can be deduced in a similar way starting again from (4.1). We summarize in table 5 the final results. In each coordinate system we obtain several equivalent expressions for the rhs of the cycle equations; the last one involves y , x or r only through $T_{\kappa_1 \kappa_2}(y/2)$, $T_{\kappa_1}(x/2)$ or $T_{\kappa_1}(r/2)$. In this last form the cycle equations in polar coordinates can be rewritten as a quadratic equation in $T_{\kappa_1}(r/2)$:

$$T_{\kappa_1}^2(r/2) - \frac{1}{\kappa_1} \frac{2}{\alpha + \alpha_0} (\alpha_1 C_{\kappa_2}(\phi) + \alpha_2 S_{\kappa_2}(\phi)) T_{\kappa_1}(r/2) + \frac{1}{\kappa_1} \frac{\alpha - \alpha_0}{\alpha + \alpha_0} = 0. \quad (4.5)$$

This equation clearly displays the universal validity, for any curved CK space, of the well known power property of Euclidean circles. Let $T_{\kappa_1}(r_1/2), T_{\kappa_1}(r_2/2)$ be the two roots of this equation, where r_1, r_2 are the distances to the origin O of the two intersection points of the cycle with the line $\phi = \text{constant}$ through O , then the product

$$T_{\kappa_1}(r_1/2) T_{\kappa_1}(r_2/2) = \frac{1}{\kappa_1} \frac{\alpha - \alpha_0}{\alpha + \alpha_0} =: \wp \quad (4.6)$$

is the same for all lines through O . As the cycle is arbitrary, this means that the same property holds for any other point. The quantity $\wp$ will be called *power of the point relative to the cycle*; in the flat limit this reduces to $r_1 r_2 = 4\wp$.

In what follows we proceed to identify the equations of the three generic types of cycles: *geodesics* (zero geodesic curvature), *equidistants* (constant distance to a geodesic) and *circles* (constant distance to a point), that are also displayed in table 5.

4.1 Geodesics

If we set $\alpha = 0$ in (4.2) then $k_g = 0$ and we obtain *geodesics*. The generic geodesic equations (when $\alpha_2 \neq 0$), which generalize the usual cartesian equations $y = \beta_0 + \beta_1 a$ ($\beta_0 = -\alpha_0/\alpha_2, \beta_1 = -\alpha_1/\alpha_2, \alpha_2 \neq 0$) reducing to them when $\kappa_1 = 0$, are:

$$T_{\kappa_1 \kappa_2}(y) = \beta_0 C_{\kappa_1}(a) + \beta_1 S_{\kappa_1}(a). \quad (4.7)$$

The case $\alpha_2 = 0$ gives the non-generic geodesics $a = a_0$. These expressions could also have been deduced by integrating the usual geodesic differential equations. In the relativistic spacetimes with $\kappa_2 = -1/c^2 < 0$, the geodesic (4.7) can be either a time-like, space-like or isotropic line according to the following conditions

$$\begin{aligned} \text{Time-like geodesic:} & \quad 1 + \kappa_1 \kappa_2 \beta_0^2 + \kappa_2 \beta_1^2 > 0 \\ \text{Space-like geodesic:} & \quad 1 + \kappa_1 \kappa_2 \beta_0^2 + \kappa_2 \beta_1^2 < 0 \\ \text{Isotropic geodesic:} & \quad 1 + \kappa_1 \kappa_2 \beta_0^2 + \kappa_2 \beta_1^2 = 0. \end{aligned} \quad (4.8)$$

Hereafter, when dealing with geodesics we will distinguish the possible types; this will be actually meaningful only whenever $\kappa_2 < 0$ and becomes irrelevant in the Riemannian spaces with $\kappa_2 > 0$, where all geodesics merge in a single type.

4.2 Equidistants

In the relativistic spacetimes, the equidistant to a given geodesic is of the same type as the geodesic, which can also be either time-like or space-like, but its equidistance radius is space-like or time-like, respectively.

The two branches of the *equidistant* to a ‘space-like’ line given by (4.7) with ‘time-like’ equidistance radius d is:

$$S_{\kappa_1}^2(d) = \kappa_2 \frac{\{S_{\kappa_1 \kappa_2}(y) - C_{\kappa_1 \kappa_2}(y)(\beta_0 C_{\kappa_1}(a) + \beta_1 S_{\kappa_1}(a))\}^2}{1 + \kappa_2 \beta_1^2 + \kappa_1 \kappa_2 \beta_0^2} \quad \text{for } \kappa_2 \neq 0. \quad (4.9)$$

These are particular cycles with equation (4.2) for $\alpha_0 = -\beta_0 \alpha_2, \alpha_1 = -\beta_1 \alpha_2, \alpha_2 \neq 0$ as given for the baseline geodesic, but with

$$\alpha = \pm \alpha_2 S_{\kappa_1}(d) (\beta_1^2 + \kappa_1 \beta_0^2 + 1/\kappa_2)^{1/2} \quad (4.10)$$

instead of $\alpha = 0$, and their geodesic curvature (4.4) reads

$$k_g = |\kappa_1 T_{\kappa_1}(d)| \quad (4.11)$$

reducing to $k_g = 0$ (geodesics) in the flat case with $\kappa_1 = 0$.

4.3 Circles

A ‘space-like’ *circle* of center (a_0, y_0) and ‘time-like’ radius ρ (see appendix B) is:

$$C_{\kappa_1}(\rho) = C_{\kappa_1\kappa_2}(y)C_{\kappa_1\kappa_2}(y_0)C_{\kappa_1}(a - a_0) + \kappa_1\kappa_2S_{\kappa_1\kappa_2}(y)S_{\kappa_1\kappa_2}(y_0) \quad (4.12)$$

and can alternatively be written in terms of versed sines (3.4):

$$V_{\kappa_1}(\rho) = C_{\kappa_1\kappa_2}(y)C_{\kappa_1\kappa_2}(y_0)V_{\kappa_1}(a - a_0) + \kappa_2V_{\kappa_1\kappa_2}(y - y_0). \quad (4.13)$$

These equations are equivalent to one in the form (4.2) for the following choice of α_i :

$$\alpha_0 = C_{\kappa_1}(a_0)C_{\kappa_1\kappa_2}(y_0) \quad \alpha_1 = \kappa_1S_{\kappa_1}(a_0)C_{\kappa_1\kappa_2}(y_0) \quad \alpha_2 = \kappa_1\kappa_2S_{\kappa_1\kappa_2}(y_0) \quad \alpha = C_{\kappa_1}(\rho). \quad (4.14)$$

Hence the geodesic curvature (4.4) for a circle reads

$$k_g = |1/T_{\kappa_1}(\rho)| \quad (4.15)$$

reducing to $k_g = 1/\rho$ in the flat case. The length $L_\psi(\rho)$ of an arc of circle and the area $\mathcal{S}_\psi(\rho)$ of a sector of circle, both of radius ρ with central angle ψ , turn out to be

$$L_\psi(\rho) = S_{\kappa_1}(\rho)\psi \quad \mathcal{S}_\psi(\rho) = V_{\kappa_1}(\rho)\psi. \quad (4.16)$$

The circle equations also give the finite form of the *distance* $s \equiv \rho$ between two points and can be obtained either by integrating the main metric $(ds)_1$ along a geodesic, or directly from the cosine theorems of the CK spaces given in [16]; this is sketched in appendix B.

To end this section, we analyze the different types of ‘space-like’ cycles obtained above which arise in the CK spaces; these can be classed according to the geodesic curvature k_g that ranges in the interval $k_g \in [0, \infty)$ and the sign of the CK constant κ_1 :

- If $\kappa_1 > 0$ ($\mathbf{S}^2$ and $\mathbf{AdS}^{1+1}$), both geodesic curvatures of ‘space-like’ equidistants (4.11) and ‘space-like’ circles (4.15) range in $k_g \in (0, \infty)$, so that such equidistants (4.9) are at once circles (4.12) and conversely; this is clear on the sphere where parallels are simultaneously equidistants to the equator and circles with center in the pole.
- If $\kappa_1 = 0$ ($\mathbf{E}^2$ and $\mathbf{M}^{1+1}$), equidistants are simply geodesics with $k_g = 0$, while circles correspond to $k_g \in (0, \infty)$.
- If $\kappa_1 < 0$ ($\mathbf{H}^2$ and $\mathbf{dS}^{1+1}$), ‘space-like’ equidistants (4.9) and circles (4.12) are *different* cycles and a third type, naturally separating them appear; these are the ‘space-like’ *horocycles*, which are the common limits of equidistants when the ‘space-like’ base geodesic goes to infinity or of ‘space-like’ circles when the center goes to infinity. In this case, equidistants correspond to the values of $k_g \in (0, \sqrt{-\kappa_1})$ while circles have $k_g \in (\sqrt{-\kappa_1}, \infty)$, with $k_g = 0$ for geodesics, $k_g = \sqrt{-\kappa_1}$ for horocycles and $k_g = \infty$ for circles of radius 0.

Equations for ‘time-like’ equidistants to a ‘time-like’ line and ‘time-like’ circles (with ‘space-like’ equidistance or radius) can also be deduced. For them an analysis similar to the above one can be performed. The classification depends on the sign of $\kappa_1\kappa_2$; for instance, in $\mathbf{H}^2$ or in $\mathbf{AdS}^{1+1}$ ($\kappa_1\kappa_2 < 0$) there are three types of ‘time-like’ cycles: equidistants to a ‘time-like’ geodesic, ‘time-like’ horocycles and ‘time-like’ circles (‘space-like’ radius). In $\mathbf{S}^2$ and in $\mathbf{dS}^{1+1}$ ($\kappa_1\kappa_2 > 0$) ‘time-like’ equidistants and ‘time-like’ circles coincide.

In table 6 we explicitly write the equations of geodesics and circles for each of the nine spaces $S^2_{[\kappa_1], \kappa_2}$ in parallel I coordinates (a complete study of Galilean lines and cycles can be found in [13]).

5 Cycle-preserving Lie groups

5.1 One-parameter subgroups of geodesic-preserving transformations

In this section we give a *direct* derivation of the group of cycle-preserving transformations in $S^2_{[\kappa_1], \kappa_2}$. We start by discussing the geodesic-preserving transformations, thus allowing to recover the CK group $SO_{\kappa_1, \kappa_2}(3)$. Consider the equations of geodesics in parallel I coordinates given in table 5. We look whether there exists a transformation within the ansatz $(a, y) \rightarrow (a'(a), y)$ which carries geodesics into geodesics. Noticing that y is assumed not to change, this requirement is equivalent to impose:

$$\beta'_0 C_{\kappa_1}(a'(a)) + \beta'_1 S_{\kappa_1}(a'(a)) \propto \beta_0 C_{\kappa_1}(a) + \beta_1 S_{\kappa_1}(a). \quad (5.1)$$

The addition properties of the trigonometric functions $C_{\kappa_1}(a), S_{\kappa_1}(a)$ [16]:

$$C_{\kappa}(x \pm y) = C_{\kappa}(x)C_{\kappa}(y) \mp \kappa S_{\kappa}(x)S_{\kappa}(y) \quad S_{\kappa}(x \pm y) = S_{\kappa}(x)C_{\kappa}(y) \pm S_{\kappa}(y)C_{\kappa}(x) \quad (5.2)$$

display directly the obvious solution $a' = a + a_0$, so that the transformation

$$(a, y) \rightarrow (a + a_0, y) \quad (5.3)$$

is indeed geodesic preserving; it is the *translation* along the geodesic l_1 of $S^2_{[\kappa_1], \kappa_2}$ generated by P_1 (see figure 1); in the kinematical language, this would be a time translation. Now we look for a geodesic-preserving transformation described by the ansatz $(x, b) \rightarrow (x, b'(b))$ in parallel II coordinates; similar arguments lead to another geodesic-preserving map:

$$(x, b) \rightarrow (x, b + b_0) \quad (5.4)$$

which is the *translation* along the geodesic l_2 generated by P_2 (a space translation). Finally, we look within the ansatz $(r, \phi) \rightarrow (r, \phi'(\phi))$ in polar coordinates; the condition of preserving geodesics singles out the transformation:

$$(r, \phi) \rightarrow (r, \phi + \phi_0) \quad (5.5)$$

which is the *rotation* around the origin of $S^2_{[\kappa_1], \kappa_2}$ with generator J_{12} (kinematically a boost). The fundamental vector fields of (5.3)–(5.5) are given by:

$$P_1 = -\partial_a \quad P_2 = -\partial_b \quad J_{12} = -\partial_{\phi}. \quad (5.6)$$

By writing them in a single coordinate system (see table 7 below) it can be shown that they close a Lie algebra, with commutators (2.1): thus we recover the CK algebra $so_{\kappa_1, \kappa_2}(3)$.

5.2 One-parameter subgroups of cycle-preserving transformations

Our purpose now is to proceed in a similar way in order to deduce the *cycle-preserving transformations* of the spaces $S^2_{[\kappa_1], \kappa_2}$ giving rise to their corresponding conformal groups. Again we start by assuming $\kappa_1 \neq 0$; at the end we will recover the $\kappa_1 \rightarrow 0$ limit.

The full CK motion group generated by $\{P_1, P_2, J_{12}\}$ is clearly cycle-preserving. Are there additional cycle-preserving transformations which are not geodesic-preserving? A

natural idea is to look for them within three ansatzes complementary to the ones previously explored, that is, $(a, y) \rightarrow (a, y'(y))$, $(x, b) \rightarrow (x'(x), b)$ and $(r, \phi) \rightarrow (r'(r), \phi)$, respectively.

Let us begin with the ansatz $(a, y) \rightarrow (a, y'(y))$ in parallel I coordinates, and require this to be cycle-preserving; taking (4.3) into account, and noticing that a is assumed not to change this is equivalent to enforce

$$\alpha' C_{-\kappa_1 \kappa_2}(y'^{\wedge}) - \alpha'_2 S_{-\kappa_1 \kappa_2}(y'^{\wedge}) \propto \alpha C_{-\kappa_1 \kappa_2}(y^{\wedge}) - \alpha_2 S_{-\kappa_1 \kappa_2}(y^{\wedge}) \quad (5.7)$$

and again the addition relations (5.2) display a solution $y'^{\wedge} = y^{\wedge} + \xi$, so that

$$(a, y) \rightarrow (a, y'(y)) \quad \Lambda_{\kappa_1 \kappa_2}(y') = \Lambda_{\kappa_1 \kappa_2}(y) + \xi \quad (5.8)$$

satisfies (5.7) and is cycle-preserving. This is a one-parametric subgroup of transformations with canonical parameter ξ ; the fundamental vector field follows using (A.12):

$$L_2 = -\partial_{y^{\wedge}} = -C_{\kappa_1 \kappa_2}(y) \partial_y. \quad (5.9)$$

Transformations generated by L_2 behave in the neighbourhood of the origin O as translations; we call L_2 the generator of Λ -translations along the geodesic l_2 of $S^2_{[\kappa_1], \kappa_2}$. On the variable $y^{\wedge} = \Lambda_{\kappa_1 \kappa_2}(y)$, L_2 generates actually ordinary translations. If one now takes into account that the inverse function of Λ_{κ} is $\Lambda_{-\kappa}$ (A.9), then (5.8) can be rewritten as

$$(a, y) \rightarrow (a, y'(y)) \quad y' = \Lambda_{-\kappa_1 \kappa_2}(\Lambda_{\kappa_1 \kappa_2}(y) + \xi). \quad (5.10)$$

Likewise, the transformation

$$(x, b) \rightarrow (x'(x), b) \quad x' = \Lambda_{-\kappa_1}(\Lambda_{\kappa_1}(x) + \zeta) \quad (5.11)$$

with canonical parameter ζ is also cycle-preserving, and its fundamental vector field is the generator of Λ -translations along the geodesic l_1 of $S^2_{[\kappa_1], \kappa_2}$:

$$L_1 = -\partial_{x^{\wedge}} = -C_{\kappa_1}(x) \partial_x. \quad (5.12)$$

There remains to search for a cycle preserving transformation within $(r, \phi) \rightarrow (r'(r), \phi)$. In this case it is better to use the cycle equation in the second alternative form given in table 5, introducing the power $\wp$ of the origin relative to the cycle (4.6), and taking into account that ϕ is assumed not to change. Thus it suffices to enforce:

$$\frac{1}{\wp'} T_{\kappa_1}(r'/2) + \frac{1}{T_{\kappa_1}(r'/2)} \propto \frac{1}{\wp} T_{\kappa_1}(r/2) + \frac{1}{T_{\kappa_1}(r/2)}. \quad (5.13)$$

Hence, the one-parametric family of transformations with canonical parameter λ :

$$T_{\kappa_1}(r'/2) = e^{\lambda} T_{\kappa_1}(r/2) \quad (5.14)$$

are cycle-preserving as they verify (5.13) with $\wp' = e^{2\lambda} \wp$; they can be interpreted as homoteties around the origin with scale factor e^{λ} . The fundamental vector field is obtained with the aid of (3.5):

$$D = -S_{\kappa_1}(r) \partial_r. \quad (5.15)$$

We remark the natural appearance of a one-parameter family of ‘dilations’ in spaces like the sphere and the hyperbolic plane, with non-zero curvature, in spite of a vague but widespread belief against this possibility.

When $\kappa_1 = 0$, the approach discussed in this paragraph only leads to the standard *dilation* $r' = e^\lambda r$ as a new transformation beyond the motions. The generators $L_1 = -C_{\kappa_1}(x) \partial_x$, $L_2 = -C_{\kappa_1 \kappa_2}(y) \partial_y$, are still meaningful in the limit $\kappa_1 = 0$ where they reduce to P_1, P_2 and do not provide any new transformation. Nevertheless, the flat conformal groups can still be obtained as some suitable limit, to be discussed below.

5.3 Involutive discrete cycle-preserving transformations

The equations (5.7) or (5.13) allow other discrete solutions not belonging to one-parametric families. These solutions correspond to *inversions in cycles* and are involutive. There are three families, each matching perfectly with the three earlier types D, L_1, L_2 . We start with the *inversions in circles*, associated to D . The discrete transformation:

$$T_{\kappa_1}(r'/2) \cdot T_{\kappa_1}(r/2) = \wp_0 \quad (5.16)$$

is also a solution of (5.13) with $\wp' = \wp = \wp_0$. The circle with center at the origin and radius ρ_0 such that $T_{\kappa_1}^2(\rho_0/2) = -\wp_0$ is invariant under this transformation, which therefore corresponds to the inversion in that circle; notice that, in this case, if $r > 0$, then $r' < 0$ and conversely, so that $\wp_0 < 0$. For any value of κ_1 , the product of two inversions in two concentric circles in the family (5.16) with constants $\wp_0, \wp'_0$ is a dilation (5.14) with the same center and scale factor $e^\lambda = \wp'_0/\wp_0$. All this is well known for the $\kappa_1 = 0$ case, where (5.16) reduces to the standard flat inversion in circles $r' = 4\wp_0/r$.

In the generic $\kappa_1 \neq 0$ case, a fully analogous situation happens for the two additional one-parameter subgroups generated by L_1, L_2 . Both Λ -translations (5.10) and (5.11) can be expressed as a product of two involutive discrete transformations, namely *inversions in equidistants*. In the $\kappa_1 = 0$ limit, however, these new inversions reduce to the ordinary reflections in geodesics (recall that L_1, L_2 also reduce in this limit to P_1, P_2).

Specifically, the rhs of the cycle equations in parallel II coordinates can be recast in the third form given in table 5. If the condition of carrying cycles into cycles is enforced, this gives an equation fully analogous to (5.13), where the rhs is replaced by:

$$\frac{1}{\kappa_1 \wp_2} \left(\frac{1 - \sqrt{\kappa_1} T_{\kappa_1}(x/2)}{1 + \sqrt{\kappa_1} T_{\kappa_1}(x/2)} \right) + \left(\frac{1 - \sqrt{\kappa_1} T_{\kappa_1}(x/2)}{1 + \sqrt{\kappa_1} T_{\kappa_1}(x/2)} \right)^{-1} \wp_2 = \frac{1}{\kappa_1} \frac{\alpha \sqrt{\kappa_1} - \alpha_1}{\alpha \sqrt{\kappa_1} + \alpha_1} \quad (5.17)$$

and the lhs is formally similar with $\alpha', \alpha'_1, x', \wp'_2$. In this form the equation allows us to read directly a discrete and involutive cycle-preserving transformation similar to (5.16):

$$\left(\frac{1 - \sqrt{\kappa_1} T_{\kappa_1}(x'/2)}{1 + \sqrt{\kappa_1} T_{\kappa_1}(x'/2)} \right) \cdot \left(\frac{1 - \sqrt{\kappa_1} T_{\kappa_1}(x/2)}{1 + \sqrt{\kappa_1} T_{\kappa_1}(x/2)} \right) = \kappa_1 \wp_2. \quad (5.18)$$

This transformation keeps invariant one branch of a certain equidistant (with parameters $\alpha_0 = \beta_0 = 0, \alpha_1 \neq 0, \alpha_2 = \beta_2 = 0, \alpha = \alpha_1 S_{\kappa_1}(d)$) to the line l_2 , and the ‘time-like’ equidistance d coincides, up to the sign, with the x coordinate which is constant along the equidistant. The quantity $\wp_2$ may well be called *power of the baseline relative to that cycle*. Again the product of two such inversions in two coaxial equidistants (with the same baseline) is a Λ -translation along the line l_1 , in total analogy with the decomposition of a dilation as the product of two inversions in concentric circles. Everything is similar for the third family corresponding to the cycle equations in parallel I coordinates, described

by similar expressions with the replacements $\alpha \rightarrow \alpha$, $\alpha_1 \rightarrow \alpha_2$, $y \rightarrow x$ and $\kappa_1 \rightarrow \kappa_1 \kappa_2$, so that $\wp_2 \rightarrow \wp_1 = \frac{1}{\kappa_1 \kappa_2} \frac{\alpha \sqrt{\kappa_1 \kappa_2 - \alpha_2}}{\alpha \sqrt{\kappa_1 \kappa_2 + \alpha_2}}$.

Thus these discrete inversions are even more basic than the one-parameter transformations generated by D, L_1, L_2 , and they display a behaviour which is completely symmetric in the generic curved case; this symmetry disappears in the flat limit.

5.4 Conformal algebras

Summing up, for $\kappa_1 \neq 0$ we have found six one-parametric subgroups of cycle-preserving transformations in $S_{[\kappa_1], \kappa_2}^2$ with generators $\{P_i, J_{12}, L_i, D\}$ ($i = 1, 2$). These close a Lie algebra. This fact can be checked by explicit computation once all the generators are written in the same coordinate system as is shown in table 7; their Lie brackets are:

$$\begin{aligned} [J_{12}, P_1] &= P_2 & [J_{12}, P_2] &= -\kappa_2 P_1 & [P_1, P_2] &= \kappa_1 J_{12} \\ [J_{12}, L_1] &= L_2 & [J_{12}, L_2] &= -\kappa_2 L_1 & [L_1, L_2] &= -\kappa_1 J_{12} \\ [D, P_i] &= L_i & [D, L_i] &= P_i & [D, J_{12}] &= 0 \\ [P_1, L_1] &= \kappa_1 D & [P_2, L_2] &= \kappa_1 \kappa_2 D & & \\ [P_1, L_2] &= 0 & [P_2, L_1] &= 0. & & \end{aligned} \quad (5.19)$$

This Lie algebra will be denoted $\text{conf}_{\kappa_1, \kappa_2}$ ($\text{m\"ob}_{\kappa_1, \kappa_2}$ would be adequate as well) and has two Casimir invariants:

$$\begin{aligned} \mathcal{C}_1 &= -\kappa_1 J_{12}^2 + \kappa_1 \kappa_2 D^2 + \kappa_2 (L_1^2 - P_1^2) + L_2^2 - P_2^2 \\ \mathcal{C}_2 &= \kappa_1 J_{12} D + L_1 P_2 - P_1 L_2. \end{aligned} \quad (5.20)$$

The basis $\{P_i, J_{12}, L_i, D\}$ shows a clear conformal *duality* or algebraic symmetric role between the generators of translations and Λ -translations: the map defined by

$$P_i \leftrightarrow L_i \quad J_{12} \leftrightarrow J_{12} \quad D \leftrightarrow D \quad (5.21)$$

interchanges the set of conformal algebras $\text{conf}_{\kappa_1, \kappa_2} \leftrightarrow \text{conf}_{-\kappa_1, \kappa_2}$.

Now we discuss the limit $\kappa_1 \rightarrow 0$. In this case both generators L_i coincide with the translation generators P_i , but as far as $\kappa_1 \neq 0$, we may take other two generators

$$G_i = \frac{1}{\kappa_1} (L_i - P_i) \quad i = 1, 2 \quad (5.22)$$

generating the so called *specific conformal transformations*. The G_i are always defined, but when $\kappa_1 = 0$ they continue to be independent of the four remaining generators P_i, J_{12}, D . The differential realization of generators $\{P_i, J_{12}, G_i, D\}$ for any value of κ_1 are given in table 7. When $\kappa_1 = 0$ and in parallel I coordinates they reproduce the known vector fields for the conformal generators of flat spaces ($\mathbf{E}^2$ or $\mathbf{M}^{1+1}$ according to the sign of κ_2):

$$\begin{aligned} P_1 &= -\partial_a & P_2 &= -\partial_y & J_{12} &= \kappa_2 y \partial_a - a \partial_y & D &= -a \partial_a - y \partial_y \\ G_1 &= \frac{1}{2}(a^2 - \kappa_2 y^2) \partial_a + a y \partial_y & G_2 &= \kappa_2 a y \partial_a - \frac{1}{2}(a^2 - \kappa_2 y^2) \partial_y. \end{aligned} \quad (5.23)$$

In the basis $\{P_i, J_{12}, G_i, D\}$ the commutation rules and Casimirs of $\text{conf}_{\kappa_1, \kappa_2}$ read

$$\begin{aligned} [J_{12}, P_1] &= P_2 & [J_{12}, P_2] &= -\kappa_2 P_1 & [P_1, P_2] &= \kappa_1 J_{12} \\ [J_{12}, G_1] &= G_2 & [J_{12}, G_2] &= -\kappa_2 G_1 & [G_1, G_2] &= 0 \\ [D, P_i] &= P_i + \kappa_1 G_i & [D, G_i] &= -G_i & [D, J_{12}] &= 0 \\ [P_1, G_1] &= D & [P_2, G_2] &= \kappa_2 D & & \\ [P_1, G_2] &= -J_{12} & [P_2, G_1] &= J_{12} & & \end{aligned} \quad (5.24)$$

$$\begin{aligned} \mathcal{C}_1 &= -J_{12}^2 + \kappa_2 D^2 + \kappa_2(P_1 G_1 + G_1 P_1) + P_2 G_2 + G_2 P_2 + \kappa_1 \kappa_2 G_1^2 + \kappa_1 G_2^2 \\ \mathcal{C}_2 &= J_{12} D + G_1 P_2 - P_1 G_2. \end{aligned} \quad (5.25)$$

This is the basis we will use in the next sections. Remarkable subalgebras of $\text{conf}_{\kappa_1, \kappa_2}$ are:

- $\{P_1, P_2, J_{12}\}$ span a CK algebra $so_{\kappa_1, \kappa_2}(3)$ (see table 1), but the set $\{P_1, P_2, J_{12}; D\}$ only close a Lie algebra (isometries plus dilations) if $\kappa_1 = 0$.
- $\{G_1, G_2, J_{12}\}$ span a *flat* CK algebra $so_{0, \kappa_2}(3)$, no matter of the value of κ_1 . The corresponding CK group has a semidirect product structure: $SO_{0, \kappa_2}(3) = ISO_{\kappa_2}(3) = T_2 \odot SO_{\kappa_2}(2)$ where $T_2 = \langle G_1, G_2 \rangle$ and $SO_{\kappa_2}(2) = \langle J_{12} \rangle$.
- $\{G_1, G_2, J_{12}, D\}$ span a Lie algebra sim_{0, κ_2} of a similitude group of a 2D flat space with the following structure: $\text{Sim}_{0, \kappa_2} = T_2 \odot (SO_{\kappa_2}(2) \otimes SO(1, 1))$, where $SO(1, 1) = \langle D \rangle$. Thus this is either the Euclidean ($\kappa_2 > 0$), Galilean ($\kappa_2 = 0$), or Poincaré ($\kappa_2 < 0$) similitude group.

Another interesting basis for $\text{conf}_{\kappa_1, \kappa_2}$ is $\{R_i, J_{12}, G_i, D\}$ where

$$R_i = P_i + \frac{1}{2} \kappa_1 G_i = \frac{1}{2} (P_i + L_i) \quad i = 1, 2. \quad (5.26)$$

The commutation relations and Casimir operators are now given by

$$\begin{aligned} [J_{12}, R_1] &= R_2 & [J_{12}, R_2] &= -\kappa_2 R_1 & [R_1, R_2] &= 0 \\ [J_{12}, G_1] &= G_2 & [J_{12}, G_2] &= -\kappa_2 G_1 & [G_1, G_2] &= 0 \\ [D, R_i] &= R_i & [D, G_i] &= -G_i & [D, J_{12}] &= 0 \\ [R_1, G_1] &= D & [R_2, G_2] &= \kappa_2 D & & \\ [R_1, G_2] &= -J_{12} & [R_2, G_1] &= J_{12} & & \end{aligned} \quad (5.27)$$

$$\begin{aligned} \mathcal{C}_1 &= -J_{12}^2 + \kappa_2 D^2 + \kappa_2(R_1 G_1 + G_1 R_1) + R_2 G_2 + G_2 R_2 \\ \mathcal{C}_2 &= J_{12} D + G_1 R_2 - R_1 G_2. \end{aligned} \quad (5.28)$$

Therefore, the coefficient κ_1 disappears from the commutators. This means that *all* spaces in the family $S_{[\kappa_1], \kappa_2}^2$ with the *same* κ_2 have isomorphic conformal algebras, no matter of the value of κ_1 . These are either:

- $so(3, 1)$ ((2+1)D de Sitter algebra) as the conformal algebra of the three 2D homogeneous Riemannian spaces with $\kappa_2 > 0$.
- $iso(2, 1)$ ((2+1)D Poincaré) for the (1+1)D non-relativistic spacetimes with $\kappa_2 = 0$.
- $so(2, 2)$ ((2+1)D anti-de Sitter) for the (1+1)D relativistic spacetimes with $\kappa_2 < 0$.

The known symmetry between P_i and G_i which exists in the flat case *does not* survive in the curved case. Note also that the vector fields of $\text{conf}_{\kappa_1, \kappa_2}$ displayed in table 7 are zero-realizations for both Casimirs. All conformal generators so far determined actually satisfy the conformal Killing equations for the metrics g_1 and g_2 ; this is checked in appendix C.

6 Conformal symmetries of Laplace/wave-type equations

As a byproduct of the conformal vector fields deduced in the previous section, we now proceed to obtain differential equations with conformal algebra symmetry.

Let us consider a 2D space with coordinates (u^1, u^2) , a differential operator $E = E(u^1, u^2, \partial_1, \partial_2)$ acting on functions $\Phi(u^1, u^2)$ defined on the space ($\partial_i \equiv \partial/\partial u^i$), and

consider the differential equation:

$$E\Phi(u^1, u^2) = 0. \quad (6.1)$$

An operator $\mathcal{O}$ is a symmetry of (6.1) if $\mathcal{O}$ transforms solutions into solutions:

$$E\mathcal{O} = \mathcal{Q}E \quad \text{or} \quad [E, \mathcal{O}] = \mathcal{Q}'E \quad (6.2)$$

where $\mathcal{Q}$ is another operator and $\mathcal{Q}' = \mathcal{Q} - \mathcal{O}$.

We now focus attention in the differential equation obtained by taking as E the Casimir $\mathcal{C}$ of the CK algebra $so_{\kappa_1, \kappa_2}(3)$ (2.2) in the space $S_{[\kappa_1], \kappa_2}^2$: $\mathcal{C}\Phi = 0$. In the three geodesic coordinate systems, such an equation turns out to be

$$\begin{aligned} & \left(\frac{\kappa_2}{C_{\kappa_1 \kappa_2}^2(y)} \partial_a^2 + \partial_y^2 - \kappa_1 \kappa_2 T_{\kappa_1 \kappa_2}(y) \partial_y \right) \Phi(a, y) = 0 \\ & \left(\kappa_2 \partial_x^2 - \kappa_1 \kappa_2 T_{\kappa_1}(x) \partial_x + \frac{1}{C_{\kappa_1}^2(x)} \partial_b^2 \right) \Phi(x, b) = 0 \\ & \left(\kappa_2 \partial_r^2 + \frac{\kappa_2}{T_{\kappa_1}(r)} \partial_r + \frac{1}{S_{\kappa_1}^2(r)} \partial_\phi^2 \right) \Phi(r, \phi) = 0. \end{aligned} \quad (6.3)$$

The conformal algebra $\text{conf}_{\kappa_1, \kappa_2}$ is a symmetry algebra of these equations, and by using table 7 the generators $\{P_i, J_{12}, G_i, D\}$ are shown to fulfil

$$\begin{aligned} [\mathcal{C}, X] &= 0 \quad X \in \{P_1, P_2, J_{12}\} \\ [\mathcal{C}, D] &= -2C_{\kappa_1}(a)C_{\kappa_1 \kappa_2}(y)\mathcal{C} = -2C_{\kappa_1}(x)C_{\kappa_1 \kappa_2}(b)\mathcal{C} = -2C_{\kappa_1}(r)\mathcal{C} \equiv -2x^0\mathcal{C} \\ [\mathcal{C}, G_1] &= 2S_{\kappa_1}(a)C_{\kappa_1 \kappa_2}(y)\mathcal{C} = 2S_{\kappa_1}(x)\mathcal{C} = 2S_{\kappa_1}(r)C_{\kappa_2}(\phi)\mathcal{C} \equiv 2x^1\mathcal{C} \\ [\mathcal{C}, G_2] &= 2\kappa_2 S_{\kappa_1 \kappa_2}(y)\mathcal{C} = 2\kappa_2 C_{\kappa_1}(x)S_{\kappa_1 \kappa_2}(b)\mathcal{C} = 2\kappa_2 S_{\kappa_1}(r)S_{\kappa_2}(\phi)\mathcal{C} \equiv 2\kappa_2 x^2\mathcal{C}. \end{aligned} \quad (6.4)$$

The operator $\mathcal{C}$ leading to the equations (6.3) is written for each specific CK space in table 8 in geodesic parallel I and polar coordinates; in the six spacetimes we introduce the notation used in tables 3 and 4. By taking into account table 8 we see:

- The usual 2D Laplace equation appears in $\mathbf{E}^2$; the corresponding non-zero curvature Laplace–Beltrami versions arise in the sphere and hyperbolic plane. All of them share the same symmetry algebra $so(3, 1)$.
- In the non-relativistic spacetimes we find an equation which does not involve time and indeed reduces to a 1D ‘Laplace’ equation. This agrees with the known absence of a true Galilean invariant wave equation and is the main reason precluding a further development of non-relativistic electromagnetic theories, where only two separate electric and magnetic essentially static limits are allowed [26, 27]; see also [28, 29] for a recent study of Newtonian electromagnetisms.
- The proper $(1 + 1)$ D wave equation is associated to $\mathbf{M}^{1+1}$ [30]; its curvature versions correspond to anti-de Sitter and de Sitter electromagnetisms in both $\mathbf{AdS}^{1+1}$ and $\mathbf{dS}^{1+1}$. These three wave-type equations have $so(2, 2)$ as their conformal symmetry algebra.

It is worth noting that this collective treatment enables a clear view of the contraction limits between these Laplace/wave-type equations together with their associated conformal symmetry algebras as shown in table 8.

7 Conformal completion of the Cayley–Klein spaces

Up to now, we have only identified the conformal group as a *local* group of transformations. The question of whether or not this group can be realized globally escapes the infinitesimal approach provided by Lie algebras. The need to suitably complete $S^2_{[\kappa_1], \kappa_2}$ with ‘additional’ points in order to have conformal transformations defined as a global *conformal group* $\text{Conf}_{\kappa_1, \kappa_2}$ arises, because the vector fields given in table 7 are, in general, not complete. An example is provided by the hyperbolic space $\mathbf{H}^2 \equiv S^2_{[-], +}$ and the finite one-parameter transformations generated by the dilation D (5.14):

$$\tanh(r'/2) = e^\lambda \tanh(r/2) \quad (7.1)$$

where e^λ is a ‘similarity factor’. A dilation with center O and factor $e^\lambda > 1$ transforms the interior of a disk of center O and radius $\rho_0 = 2 \operatorname{arctanh}(e^{-\lambda})$ into the *entire* hyperbolic space, because $\rho'_0 = \infty$. What about the images of points *outside* this disk? If one wants to have a global group, the images of these points outside the disk should be added to the ‘ordinary’ points.

The same conclusion follows from a direct analysis of the discrete inversions, expressed in terms of the functions $T_{\kappa_1}(r/2), T_{\kappa_1}(x/2), T_{\kappa_1 \kappa_2}(y/2)$, so that the problem of adding new points is related with the properties of the function $c \rightarrow T_\kappa(c/2)$. This leads us to a study of the Lambda function (see appendix A) and shows that a suitable extension of this function (mapping in a one-to-one way the *complete* elliptic line into *two copies* of the hyperbolic line glued by their final points) is required and suffices to deal with this problem algebraically.

We will however not follow here the details of this path, and to accomplish the conformal completion of the spaces $S^2_{[\kappa_1], \kappa_2}$ we sketch in this section an alternative approach which leads to the same results. This procedure extends to any value of curvature and signature the familiar linealization of the conformal flat Euclidean or Minkowskian groups.

From the group theoretical viewpoint, the conformal completion of the CK space $S^2_{[\kappa_1], \kappa_2}$ is defined as a homogeneous space $\text{Conf}_{\kappa_1, \kappa_2} / \text{Sim}_{0, \kappa_2}$, where $\text{Conf}_{\kappa_1, \kappa_2}$ is a matrix Lie group defined below and Sim_{0, κ_2} is the subgroup of $\text{Conf}_{\kappa_1, \kappa_2}$ generated by $\{J_{12}, G_1, G_2, D\}$, isomorphic to the similitude group of the flat CK space $S^2_{[0], \kappa_2}$. These groups are determined through the Lie algebras $\text{conf}_{\kappa_1, \kappa_2}$ and sim_{0, κ_2} by means of a 4D matrix realization which allows us to *define* $\text{Conf}_{\kappa_1, \kappa_2}$ as a group of *linear* transformations in a 4D *conformal ambient space*. Therefore, while the action of the CK group $SO_{\kappa_1, \kappa_2}(3)$ on the 2D spaces $S^2_{[\kappa_1], \kappa_2}$ can be linearized in an ambient space $\mathbb{R}^3$ with one extra dimension, the conformal group $\text{Conf}_{\kappa_1, \kappa_2}$ is realized as a matrix group acting as globally defined linear transformations in a ‘conformal ambient space’ $\mathbb{R}^4$, with *two* extra dimensions. To establish these ideas we first study the completion of the 1D spaces where κ_1 cannot be interpreted as the gaussian curvature, as the three 1D spaces are flat; there are however differences better described after the values of κ_1 .

7.1 One-dimensional conformal spaces

The 1D CK spaces $S^1_{[\kappa_1]}$ are the circle $\mathbf{S}^1 \equiv S^1_{[+]}$, the Euclidean line $\mathbf{E}^1 \equiv S^1_{[0]} \equiv \mathbb{R}$ and the hyperbolic line $\mathbf{H}^1 \equiv S^1_{[-]}$ (which topologically can be identified with the open segment $(-1, +1)$); the corresponding CK group $SO_{\kappa_1}(2) = \langle P_1 \rangle$ is isomorphic, in this order, to

$SO(2)$, $ISO(1) \equiv \mathbb{R}$, and $SO(1,1)$. The conformal algebra conf_{κ_1} has three generators $\{P_1, G_1, D\}$ verifying the following commutation relations:

$$[D, P_1] = P_1 + \kappa_1 G_1 \quad [D, G_1] = -G_1 \quad [P_1, G_1] = D. \quad (7.2)$$

The differential realization in terms of a single coordinate a is given by:

$$P_1 = -\partial_a \quad G_1 = V_{\kappa_1}(a) \partial_a \quad D = -S_{\kappa_1}(a) \partial_a. \quad (7.3)$$

In order to describe the conformal completion of the three spaces $S^1_{[\kappa_1]}$ we consider a 3×3 real matrix representation of conf_{κ_1} obtained through the isomorphism $\text{conf}_{\kappa_1} \simeq \text{so}(2,1)$ which can directly be seen in terms of the alternative generators $\{R_1, G_1, D\}$:

$$[D, R_1] = R_1 \quad [D, G_1] = -G_1 \quad [R_1, G_1] = D. \quad (7.4)$$

Starting from the realization of $\text{so}(2,1)$ in a pseudo-orthogonal basis $\{\Omega_{01}, \Omega_{02}, \Omega_{12}\}$ given by (3.1) with the two labels equal to -1 :

$$\Omega_{01} = e_{01} + e_{10} \quad \Omega_{02} = -e_{02} + e_{20} \quad \Omega_{12} = e_{12} + e_{21} \quad (7.5)$$

we find the matrices for the generators $\{R_1, G_1, D\}$:

$$R_1 = \frac{1}{2\ell}(\Omega_{02} - \Omega_{12}) \quad G_1 = -\ell(\Omega_{02} + \Omega_{12}) \quad D = \Omega_{01} \quad (7.6)$$

where κ_1 does not appear and ℓ is a non-zero constant with dimensions of length, which is required by dimensional reasons (R_1 and G_1 must be dimensionally inverse of each other after (7.4); see [31]). Changing back to the basis $\{P_1, G_1, D\}$ by means of (5.26), we obtain (remark that the product $\kappa_1 \ell^2$ is dimensionless):

$$P_1 = \frac{1}{2\ell} \begin{pmatrix} 0 & 0 & -1 - \kappa_1 \ell^2 \\ 0 & 0 & -1 + \kappa_1 \ell^2 \\ 1 + \kappa_1 \ell^2 & -1 + \kappa_1 \ell^2 & 0 \end{pmatrix} \\ G_1 = \ell \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & -1 \\ -1 & -1 & 0 \end{pmatrix} \quad D = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (7.7)$$

Exponentiation gives rise to the following one-parametric subgroups of Conf_{κ_1} :

$$e^{\mu P_1} = \begin{pmatrix} 1 - \frac{1}{4\ell^2}(1 + \kappa_1 \ell^2)^2 V_{\kappa_1}(\mu) & \frac{1}{4\ell^2}(1 - \kappa_1^2 \ell^4) V_{\kappa_1}(\mu) & -\frac{1}{2\ell}(1 + \kappa_1 \ell^2) S_{\kappa_1}(\mu) \\ -\frac{1}{4\ell^2}(1 - \kappa_1^2 \ell^4) V_{\kappa_1}(\mu) & 1 + \frac{1}{4\ell^2}(1 - \kappa_1 \ell^2)^2 V_{\kappa_1}(\mu) & -\frac{1}{2\ell}(1 - \kappa_1 \ell^2) S_{\kappa_1}(\mu) \\ \frac{1}{2\ell}(1 + \kappa_1 \ell^2) S_{\kappa_1}(\mu) & -\frac{1}{2\ell}(1 - \kappa_1 \ell^2) S_{\kappa_1}(\mu) & C_{\kappa_1}(\mu) \end{pmatrix} \\ e^{\nu G_1} = \begin{pmatrix} 1 - \frac{1}{2}\nu^2 \ell^2 & -\frac{1}{2}\nu^2 \ell^2 & \nu \ell \\ \frac{1}{2}\nu^2 \ell^2 & 1 + \frac{1}{2}\nu^2 \ell^2 & -\nu \ell \\ -\nu \ell & -\nu \ell & 1 \end{pmatrix} \quad e^{\xi D} = \begin{pmatrix} \cosh \xi & \sinh \xi & 0 \\ \sinh \xi & \cosh \xi & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (7.8)$$

In this representation the conformal group $\text{Conf}_{\kappa_1} \simeq SO(2,1)$ acts in a ‘conformal linear ambient space’ $\mathbb{R}^3 = (s^+, s^-, s^1)$ and one may easily check that this action is as a

group of isometries of a bilinear form $\Upsilon = \text{diag}(1, -1, 1)$, that is, any element $X \in \text{Conf}_{\kappa_1}$ fulfils

$$X^T \Upsilon X = \Upsilon. \quad (7.9)$$

Therefore the action of Conf_{κ_1} preserves the quadratic form $(s^+)^2 - (s^-)^2 + (s^1)^2$. The one-parameter subgroup $\exp(\nu G_1)$ turns out to be the isotropy subgroup of the point $\mathcal{O} = (1, -1, 0)$, while $\exp(\xi D)$ transforms this point into $(e^{-\xi}, -e^{-\xi}, 0)$. Thus the subgroup Sim_0 , generated by $\{G_1, D\}$, coincides with the isotropy subgroup of the *ray* of $\mathcal{O}$. The orbit of $\mathcal{O}$ under Conf_{κ_1} is henceforth contained in the *cone* Γ_0 given by

$$\Gamma_0 \equiv (s^+)^2 - (s^-)^2 + (s^1)^2 = 0. \quad (7.10)$$

In this way we find an explicit model of the conformal space as contained in the *projective cone* $\Gamma_0/\propto$ obtained from Γ_0 modulo the proportionality relation in the ambient space; more precisely, as the orbit of the ray of $\mathcal{O}$ in $\Gamma_0/\propto$, and Conf_{κ_1} acts transitively on this orbit. As a sideline remark, two possible notions of ray can be used here, allowing proportionality with any non-zero factor or with positive factors; they lead to two related conformal spaces, the latter being a twofold covering of the former. We shall stick to the relation given as proportionality with a non-zero factor.

Since the cone has no direction with $s^- = 0$, the projective cone, that is, the set of rays in Γ_0 can directly be identified with the section $s^- = -1$ of Γ_0 . Each point in the cone (s^+, s^-, s^1) is represented by a point $(\tilde{s}^+, \tilde{s}^1)$ in this section according to $(\tilde{s}^+ = -s^+/s^-, \tilde{s}^1 = -s^1/s^1)$, and fulfilling

$$(\tilde{s}^+)^2 + (\tilde{s}^1)^2 = 1. \quad (7.11)$$

Consequently, the 1D completed conformal space

$$C_{[\kappa_1]}^1 \equiv \text{Conf}_{\kappa_1}/\text{Sim}_0 = SO(2, 1)/(T_1 \odot SO(1, 1)) \quad \text{Sim}_0 = \langle G_1, D \rangle \quad (7.12)$$

can be identified with a *circle* $\mathbf{S}^1$:

$$C_{[\kappa_1]}^1 \equiv \Gamma_0/\propto \equiv (\tilde{s}^+)^2 + (\tilde{s}^1)^2 = 1 \longleftrightarrow \mathbf{S}^1. \quad (7.13)$$

Note that $C_{[\kappa_1]}^1$ is indeed independent of κ_1 , but we maintain reference to κ_1 to remind the original space. The natural parametric description of $C_{[\kappa_1]}^1 \equiv \mathbf{S}^1$ is in terms of a single coordinate $A \in [-\pi, \pi]$ with π and $-\pi$ identified:

$$\tilde{s}^+ = \cos A \quad \tilde{s}^1 = \sin A. \quad (7.14)$$

7.1.1 Conformal embedding $S_{[\kappa_1]}^1 \mapsto C_{[\kappa_1]}^1$

The complete CK space $S_{[\kappa_1]}^1$ is naturally embedded in $C_{[\kappa_1]}^1$; to describe the embedding the natural procedure is to choose $\mathcal{O}$ as an *origin* point, to identify $S_{[\kappa_1]}^1$ to the orbit of $\mathcal{O}$ in $\mathbf{S}^1$ under the subgroup $\exp(\mu P_1)$ that moves $\mathcal{O}$ in the cone Γ_0 , and after to consider the projective identification (7.13) in the conformal space $\mathbf{S}^1$. The action of $\exp(\mu P_1)$ on $\mathcal{O}$ provides a parametrization of the orbit of this point in the conformal ambient space $\mathbb{R}^3$

in terms of a single coordinate a :

$$\begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} \in \mathbf{S}^1 \rightarrow \begin{pmatrix} s^+ \\ s^- \\ s^1 \end{pmatrix} = e^{aP_1} \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 - \frac{1}{2\ell^2}(1 + \kappa_1\ell^2)V_{\kappa_1}(a) \\ -1 - \frac{1}{2\ell^2}(1 - \kappa_1\ell^2)V_{\kappa_1}(a) \\ S_{\kappa_1}(a)/\ell \end{pmatrix} \in \Gamma_0. \quad (7.15)$$

Within the identification (7.13) we should put the image of $(1, -1, 0)$ back to the circle determined by the intersection of the cone with the plane $s^- = -1$. Hence the explicit expression for the parametrization $(\tilde{s}^+, \tilde{s}^1)$ of the conformal space is:

$$\tilde{s}^+ = \frac{\ell^2 - T_{\kappa_1}^2(a/2)}{\ell^2 + T_{\kappa_1}^2(a/2)} \quad \tilde{s}^1 = \frac{2\ell T_{\kappa_1}(a/2)}{\ell^2 + T_{\kappa_1}^2(a/2)}. \quad (7.16)$$

By comparing with (7.14) we find the universal description of the embedding as:

$$a \in S_{[\kappa_1]}^1 \mapsto A \in C_{[\kappa_1]}^1 \quad \frac{1}{\ell^2} T_{\kappa_1}^2(a/2) = \tan^2(A/2). \quad (7.17)$$

This embedding, obtained through a group theoretical argument, has a very simple geometrical description as a *stereographic projection*. The initial CK space $S_{[\kappa_1]}^1$ lives in an ambient space $\mathbb{R}^2 = (x^0, x^1/\ell)$, with equation $(x^0)^2 + \kappa_1\ell^2(x^1/\ell)^2 = 1$, which can be considered as the CK space $S_{[\kappa_1\ell^2]}^1$. The conformal compactification $C_{[\kappa_1]}^1$ is identified with the circle $\mathbf{S}^1$ (7.13) in an ambient space $\mathbb{R}^2 = (\tilde{s}^+, \tilde{s}^1)$. Identify both ambient spaces as $(\tilde{s}^+ \leftrightarrow x^0, \tilde{s}^1 \leftrightarrow x^1/\ell)$; in this common space, the embedding of $S_{[\kappa_1]}^1$ into $C_{[\kappa_1]}^1$ is simply the stereographic projection with pole $\mathcal{P} = (-1, 0)$ of $S_{[\kappa_1\ell^2]}^1$ into $\mathbf{S}^1$, that is,

$$(x^0 + 1, x^1/\ell) = \mu(\tilde{s}^+ + 1, \tilde{s}^1) \quad (7.18)$$

where μ is a real factor which can easily be derived and reads:

$$\mu = \frac{2}{(1 + \tilde{s}^+) + \kappa_1\ell^2(1 - \tilde{s}^+)} = \frac{x^0 + 1}{2} + \frac{1 - x^0}{2\kappa_1\ell^2}. \quad (7.19)$$

Hence the stereographic projection equations turn out to be

$$\begin{aligned} \tilde{s}^+ &= \frac{\ell^2 - \frac{1}{\kappa_1} \frac{1-x^0}{1+x^0}}{\ell^2 + \frac{1}{\kappa_1} \frac{1-x^0}{1+x^0}} & x^0 &= \frac{(1 + \tilde{s}^+) - \kappa_1\ell^2(1 - \tilde{s}^+)}{(1 + \tilde{s}^+) + \kappa_1\ell^2(1 - \tilde{s}^+)} \\ \tilde{s}^1 &= \frac{2\ell \frac{1}{1+x^0}}{\ell^2 + \frac{1}{\kappa_1} \frac{1-x^0}{1+x^0}} x^1 & x^1 &= \frac{2\ell}{(1 + \tilde{s}^+) + \kappa_1\ell^2(1 - \tilde{s}^+)} \tilde{s}^1. \end{aligned} \quad (7.20)$$

By substituting $x^0 = C_{\kappa_1}(a)$ and $x^1 = S_{\kappa_1}(a)$ in (7.20) we recover (7.16). The three particular projections are represented in figure 2.

The embedding has a non-canonical character, as it depends on the choice of ℓ (or on the arbitrary choice $s^- = -1$ in (7.10)). Only the *sign* of κ_1 matters, as for any fixed κ_1 , a suitable choice of ℓ can reduce the dimensionless product $\kappa_1\ell^2$ to either 1, 0, -1. This assumed, parametrization (7.16) encompasses the three 1D conformal spaces:

$$\begin{aligned} \kappa_1\ell^2 = 1 & \quad \text{Elliptic } C_{[+]}^1 \equiv \mathbf{S}^1 : & \tilde{s}^+ &= \cos(a/\ell) & \tilde{s}^1 &= \sin(a/\ell) \\ \kappa_1\ell^2 = 0 & \quad \text{Euclidean } C_{[0]}^1 \equiv \mathbf{S}^1 : & \tilde{s}^+ &= \frac{4 - (a/\ell)^2}{4 + (a/\ell)^2} & \tilde{s}^1 &= \frac{4(a/\ell)}{4 + (a/\ell)^2} \\ \kappa_1\ell^2 = -1 & \quad \text{Hyperbolic } C_{[-]}^1 \equiv \mathbf{S}^1 : & \tilde{s}^+ &= \frac{1}{\cosh(a/\ell)} & \tilde{s}^1 &= \tanh(a/\ell) \end{aligned}$$




Figure 2: 2D visualization of the three conformal embeddings $S^1_{[\kappa_1]} \mapsto C^1_{[\kappa_1]} \equiv \mathbf{S}^1$ with coordinates $(\tilde{s}^+ \leftrightarrow x^0, \tilde{s}^1 \leftrightarrow x^1/\ell)$. The point $Q = (x^0, x^1/\ell)$ belongs to the CK space $S^1_{[\kappa_1 \ell^2]}$, while $\tilde{Q} = (\tilde{s}^+, \tilde{s}^1)$ is its stereographic projection with pole $\mathcal{P} = (-1, 0)$ in the conformal compactification $\mathbf{S}^1$; the point $O \equiv \mathcal{O} = (1, 0)$ is the origin in the spaces. The point $\tilde{R}$ is obtained through stereographic projection of R that belongs to another copy of $\mathbf{H}^1$; hence $\tilde{R}$ is a new point to be added in the conformal completion $\mathbf{S}^1$.

where we have used:

$$C_{\kappa_1}(a) = C_{\kappa_1 \ell^2}(a/\ell) \quad S_{\kappa_1}(a) = \ell S_{\kappa_1 \ell^2}(a/\ell) \quad T_{\kappa_1}(a) = \ell T_{\kappa_1 \ell^2}(a/\ell). \quad (7.21)$$

While the conformal completion of either the elliptic, Euclidean or hyperbolic line is the same circle $\mathbf{S}^1$, there are some differences between the three cases. For $\kappa_1 > 0$ the image of the whole elliptic space $S^1_{[+]}$ fills in completely the conformal space $C^1_{[+]} \equiv \mathbf{S}^1$. When $\kappa_1 < 0$ the image of the whole hyperbolic line $S^1_{[-]} \equiv \mathbf{H}^1$ does only fill a proper subinterval $\mathcal{I}_\ell = (-2 \arctan(1/\ell\sqrt{-\kappa_1}), 2 \arctan(1/\ell\sqrt{-\kappa_1})) \subset C^1_{[-]} \equiv \mathbf{S}^1$; this is clear from stereographic projection (see figure 2). Due to the arbitrariness of ℓ , the position of the images in the conformal embedding of the two final points of $\mathbf{H}^1$ is conventional (e.g. for $\kappa_1 \ell^2 = -1$, $\mathcal{I}_\ell = (-\frac{\pi}{2}, \frac{\pi}{2})$), but the fact that the completed conformal hyperbolic space $C^1_{[-]}$ contains an infinity of *new* points not already present in $\mathbf{H}^1$ is *not* conventional. These new points, filling in a subinterval complementary to $\mathcal{I}_\ell$ and centered in $A = \pi$, can be interpreted as another copy of a hyperbolic line, which is glued to the original one by their final points which are identified. In the limit $\kappa_1 \rightarrow 0$, this new copy collapses to a single point since $\mathcal{I}_\ell \rightarrow (-\pi, \pi)$. Hence only the point $A = \pi \equiv -\pi$ of $C^1_{[0]} \equiv \mathbf{S}^1$ does not appear as image of a proper point of the Euclidean line $S^1_{[0]} \equiv \mathbf{E}^1$ under the embedding (7.17); such a point coincides with the stereographic projection pole $\mathcal{P}$. Therefore one has to add no points to $\mathbf{S}^1$, a single point to $\mathbf{E}^1$ and an infinity of points to $\mathbf{H}^1$ (another copy of $\mathbf{H}^1$ indeed) in order to obtain their conformal completion $C^1_{[\kappa_1]} \equiv \mathbf{S}^1$.

7.2 Two-dimensional conformal spaces

We now follow the same steps described in the previous subsection for the 2D case. The generators $\{P_i, J_{12}, G_i, D\}$ of the conformal algebra $\text{conf}_{\kappa_1, \kappa_2}$ of the 2D spaces $S^2_{[\kappa_1], \kappa_2}$ can be regarded as generators of linear transformations in a ‘conformal ambient space’

$\mathbb{R}^4 = (s^+, s^-, s^1, s^2)$ by means of the following 4×4 matrix representation of (5.24):

$$\begin{aligned}
P_1 &= \frac{1}{2\ell} \begin{pmatrix} 0 & 0 & -1 - \kappa_1 \ell^2 & 0 \\ 0 & 0 & -1 + \kappa_1 \ell^2 & 0 \\ 1 + \kappa_1 \ell^2 & -1 + \kappa_1 \ell^2 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} & J_{12} &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\kappa_2 \\ 0 & 0 & 1 & 0 \end{pmatrix} \\
P_2 &= \frac{1}{2\ell} \begin{pmatrix} 0 & 0 & 0 & -\kappa_2(1 + \kappa_1 \ell^2) \\ 0 & 0 & 0 & -\kappa_2(1 - \kappa_1 \ell^2) \\ 0 & 0 & 0 & 0 \\ 1 + \kappa_1 \ell^2 & -1 + \kappa_1 \ell^2 & 0 & 0 \end{pmatrix} & D &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \\
G_1 &= \ell \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 \\ -1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} & G_2 &= \ell \begin{pmatrix} 0 & 0 & 0 & \kappa_2 \\ 0 & 0 & 0 & -\kappa_2 \\ 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 \end{pmatrix} \tag{7.22}
\end{aligned}$$

which follows [32] as before from the identification of the conformal algebras to either $so(3, 1)$, $iso(2, 1)$ or $so(2, 2)$ according to the sign of κ_2 . The corresponding one-parametric subgroups of $\text{Conf}_{\kappa_1, \kappa_2}$ turn out to be:

$$\begin{aligned}
e^{\mu_1 P_1} &= \begin{pmatrix} 1 - \frac{(1+\kappa_1 \ell^2)^2}{4\ell^2} V_{\kappa_1}(\mu_1) & \frac{(1-\kappa_1^2 \ell^4)}{4\ell^2} V_{\kappa_1}(\mu_1) & -\frac{(1+\kappa_1 \ell^2)}{2\ell} S_{\kappa_1}(\mu_1) & 0 \\ -\frac{(1-\kappa_1^2 \ell^4)}{4\ell^2} V_{\kappa_1}(\mu_1) & 1 + \frac{(1-\kappa_1 \ell^2)^2}{4\ell^2} V_{\kappa_1}(\mu_1) & -\frac{(1-\kappa_1 \ell^2)}{2\ell} S_{\kappa_1}(\mu_1) & 0 \\ \frac{(1+\kappa_1 \ell^2)}{2\ell} S_{\kappa_1}(\mu_1) & -\frac{(1-\kappa_1 \ell^2)}{2\ell} S_{\kappa_1}(\mu_1) & C_{\kappa_1}(\mu_1) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\
e^{\mu_2 P_2} &= \begin{pmatrix} 1 - \frac{\kappa_2(1+\kappa_1 \ell^2)^2}{4\ell^2} V_{\kappa_1 \kappa_2}(\mu_2) & \frac{\kappa_2(1-\kappa_1^2 \ell^4)}{4\ell^2} V_{\kappa_1 \kappa_2}(\mu_2) & 0 & -\frac{\kappa_2(1+\kappa_1 \ell^2)}{2\ell} S_{\kappa_1 \kappa_2}(\mu_2) \\ -\frac{\kappa_2(1-\kappa_1^2 \ell^4)}{4\ell^2} V_{\kappa_1 \kappa_2}(\mu_2) & 1 + \frac{\kappa_2(1-\kappa_1 \ell^2)^2}{4\ell^2} V_{\kappa_1 \kappa_2}(\mu_2) & 0 & -\frac{\kappa_2(1-\kappa_1 \ell^2)}{2\ell} S_{\kappa_1 \kappa_2}(\mu_2) \\ 0 & 0 & 1 & 0 \\ \frac{(1+\kappa_1 \ell^2)}{2\ell} S_{\kappa_1 \kappa_2}(\mu_2) & -\frac{(1-\kappa_1 \ell^2)}{2\ell} S_{\kappa_1 \kappa_2}(\mu_2) & 0 & C_{\kappa_1 \kappa_2}(\mu_2) \end{pmatrix} \\
e^{\nu_1 G_1} &= \begin{pmatrix} 1 - \frac{1}{2} \nu_1^2 \ell^2 & -\frac{1}{2} \nu_1^2 \ell^2 & \nu_1 \ell & 0 \\ \frac{1}{2} \nu_1^2 \ell^2 & 1 + \frac{1}{2} \nu_1^2 \ell^2 & -\nu_1 \ell & 0 \\ -\nu_1 \ell & -\nu_1 \ell & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} & e^{\psi J_{12}} &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & C_{\kappa_2}(\psi) & -\kappa_2 S_{\kappa_2}(\psi) \\ 0 & 0 & S_{\kappa_2}(\psi) & C_{\kappa_2}(\psi) \end{pmatrix} \\
e^{\nu_2 G_2} &= \begin{pmatrix} 1 - \frac{1}{2} \kappa_2 \nu_2^2 \ell^2 & -\frac{1}{2} \kappa_2 \nu_2^2 \ell^2 & 0 & \kappa_2 \nu_2 \ell \\ \frac{1}{2} \kappa_2 \nu_2^2 \ell^2 & 1 + \frac{1}{2} \kappa_2 \nu_2^2 \ell^2 & 0 & -\kappa_2 \nu_2 \ell \\ 0 & 0 & 1 & 0 \\ -\nu_2 \ell & -\nu_2 \ell & 0 & 1 \end{pmatrix} & e^{\xi D} &= \begin{pmatrix} \cosh \xi & \sinh \xi & 0 & 0 \\ \sinh \xi & \cosh \xi & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \tag{7.23}
\end{aligned}$$

The conformal group $\text{Conf}_{\kappa_1, \kappa_2}$ acts as the isometry group of the bilinear form

$$\Upsilon = \text{diag}(1, -1, 1, \kappa_2) \tag{7.24}$$

verifying the relation (7.9). Thus $\text{Conf}_{\kappa_1, \kappa_2}$ preserves the quadratic form $(s^+)^2 - (s^-)^2 + (s^1)^2 + \kappa_2 (s^2)^2$. We consider the cone Γ_0 defined by:

$$\Gamma_0 \equiv (s^+)^2 - (s^-)^2 + (s^1)^2 + \kappa_2 (s^2)^2 = 0. \tag{7.25}$$

The subgroup $\langle J_{12}, G_1, G_2 \rangle$ leaves invariant the *origin* point $\mathcal{O} = (1, -1, 0, 0) \in \Gamma_0$, while the dilation subgroup transforms $\mathcal{O}$ into $(e^{-\xi}, -e^{-\xi}, 0, 0)$. Therefore $\text{Sim}_{0, \kappa_2} = \langle J_{12}, G_1, G_2, D \rangle$ is the isotropy subgroup of the *ray* of $\mathcal{O}$; this is isomorphic to the similitude group of a flat space as commented in section 5.4. The two remaining subgroups generated by P_1 and P_2 move the ray of $\mathcal{O}$.

The 2D completed conformal space $C_{[\kappa_1], \kappa_2}^2$ is defined as the homogeneous space

$$\begin{aligned} C_{[\kappa_1], \kappa_2}^2 &\equiv \text{Conf}_{\kappa_1, \kappa_2} / \text{Sim}_{0, \kappa_2} & \text{Sim}_{0, \kappa_2} &= T_2 \odot (SO_{\kappa_2}(2) \otimes SO(1, 1)) \\ T_2 = \langle G_1, G_2 \rangle & & SO_{\kappa_2}(2) &= \langle J_{12} \rangle & SO(1, 1) &= \langle D \rangle \end{aligned} \quad (7.26)$$

and can again be identified with the orbit of the ray of $\mathcal{O}$ under the linear action of $\text{Conf}_{\kappa_1, \kappa_2}$ in $\mathbb{R}^4$. Now the situation is a bit more complicated and more interesting than in the previous 1D case, because when $\kappa_2 < 0$ the cone will contain directions with $s^- = 0$, and the direct identification with the section $s^- = -1$ of Γ_0 cannot be longer made. Consider first the rays for which $s^- \neq 0$; each such a ray determines a unique and well defined point in the section of Γ_0 by $s^- = -1$. The natural coordinates $(\tilde{s}^+, \tilde{s}^1, \tilde{s}^2)$ on this section are defined by $\tilde{s}^i = -s^i/s^-$ and verify

$$(\tilde{s}^+)^2 + (\tilde{s}^1)^2 + \kappa_2(\tilde{s}^2)^2 = 1 \longleftrightarrow S_{[+], \kappa_2}^2 \quad (7.27)$$

displaying the possible identification of the set of these rays with the orbit of $\mathcal{O}$ in a CK ‘sphere’ (3.8), hence to a CK space $S_{[+], \kappa_2}^2$. If $\kappa_2 > 0$ the cone cannot contain directions with $s^- = 0$. In this way we find that

- The three 2D *Riemannian spaces* with $\kappa_2 > 0$ and *any* κ_1 have as conformal completion the CK space $S_{[+], +}^2$, that is, the ordinary *sphere* $\mathbf{S}^2$:

$$C_{[\kappa_1], +}^2 \equiv \text{Conf}_{\kappa_1, +} / \text{Sim}_{0, +} \equiv SO(3, 1) / (T_2 \odot (SO(2) \otimes SO(1, 1))) \equiv S_{[+], +}^2 \equiv \mathbf{S}^2.$$

The situation is a bit more complicated when $\kappa_2 \leq 0$, as in these cases the cone Γ_0 will always contain rays with $s^- = 0$. Points in the conformal ambient space with $s^- = 0$ will have a ray with no proper intersection with the former section, yet they will appear as points at infinity in (7.27). Further, the fact that we are dealing with rays imply that these points should be identified through the ordinary antipodal identification in $\mathbb{R}^4$, which when $s^- = 0$ translates into antipodal identification of these points at infinity in the section $(\tilde{s}^+, \tilde{s}^- = -1, \tilde{s}^1, \tilde{s}^2)$ of the ambient space. Thus we obtain the result:

$$C_{[\kappa_1], \kappa_2}^2 = \tilde{S}_{[+], \kappa_2}^2 := S_{[+], \kappa_2}^2 \cup \{\text{points at infinity in } S_{[+], \kappa_2}^2 \text{ with antipodal identification}\}.$$

Of course, this description is still valid when $\kappa_2 > 0$, but then there are no points at infinity in $S_{[+], \kappa_2}^2$, so that $\tilde{S}_{[+], +}^2 = S_{[+], +}^2$. The space $C_{[\kappa_1], \kappa_2}^2$ so obtained is always compact (hence the name of conformal compactification). We now consider the two remaining possibilities $\kappa_2 = 0$ and $\kappa_2 < 0$ (see table 1):

- The three $(1 + 1)$ D *non-relativistic spacetimes* with $\kappa_2 = 0$ and *any* κ_1 have as their conformal compactification

$$C_{[\kappa_1], 0}^2 \equiv \text{Conf}_{\kappa_1, 0} / \text{Sim}_{0, 0} \equiv ISO(2, 1) / (T_2 \odot (ISO(1) \otimes SO(1, 1))) \equiv \tilde{S}_{[+], 0}^2 \equiv \widetilde{\mathbf{NH}}_+^{1+1}$$

obtained from the *oscillating NH spacetime* through antipodal identification of points at infinity in $\mathbf{NH}_+^{1+1}$.

- The three $(1+1)$ D *relativistic spacetimes* with $\kappa_2 < 0$ and *any* κ_1 have as their conformal compactification

$$C_{[\kappa_1],-}^2 \equiv \text{Conf}_{\kappa_1,-}/\text{Sim}_{0,-} \equiv SO(2,2)/(T_2 \odot (SO(1,1) \otimes SO(1,1))) \equiv \tilde{S}_{[+],-}^2 \equiv \widetilde{\text{AdS}}^{1+1}$$

obtained from the *anti-de Sitter spacetime* through antipodal identification of points at infinity in AdS^{1+1} .

7.2.1 Conformal embedding $S_{[\kappa_1],\kappa_2}^2 \mapsto C_{[\kappa_1],\kappa_2}^2$

Consider a point Q in $S_{[\kappa_1],\kappa_2}^2$ with coordinates (a, y) , (x, b) , (r, ϕ) , and Weierstrass coordinates (x^0, x^1, x^2) . Its image under the conformal embedding is the point $\tilde{Q}$ obtained from $\mathcal{O} = (1, -1, 0, 0) \in \Gamma_0$ under the pairs of subgroups in (7.23) naturally associated to the three types of coordinates, and the *same* canonical parameters. By following the pattern of section 3.3 we obtain parametrizations for $\tilde{Q} = (s^+, s^-, s^1, s^2) \in \Gamma_0$ (as in (7.15)) in either parallel I, parallel II or polar coordinates of the initial space $S_{[\kappa_1],\kappa_2}^2$. Next the embedding into $\tilde{S}_{[+],\kappa_2}^2$ is described explicitly by $(\tilde{s}^+, \tilde{s}^1, \tilde{s}^2)$ with $\tilde{s}^i = -s^i/s^-$. The final results are displayed in table 9; notice the simple expressions that arise in polar coordinates.

As in the 1D case, this algebraic construction allows a very neat geometrical interpretation as a *stereographic projection*. Proceed as in the 1D case, and identify the linear ambient space $\mathbb{R}^3 = (x^0, x^1/\ell, x^2/\ell)$ where the initial CK space $S_{[\kappa_1],\kappa_2}^2$ can be realized as $S_{[\kappa_1\ell^2],\kappa_2}^2$ to the space $\mathbb{R}^3 = (\tilde{s}^+, \tilde{s}^1, \tilde{s}^2)$ where $S_{[+],\kappa_2}^2$ lives. In this common space $(\tilde{s}^+ \leftrightarrow x^0, \tilde{s}^1 \leftrightarrow x^1/\ell, \tilde{s}^2 \leftrightarrow x^2/\ell)$ the embedding of $S_{[\kappa_1],\kappa_2}^2$ into $C_{[\kappa_1],\kappa_2}^2$ is simply a stereographic projection of $S_{[\kappa_1\ell^2],\kappa_2}^2$ into $S_{[+],\kappa_2}^2$ with pole $\mathcal{P} = (-1, 0, 0)$. Its corresponding equations are again given by (7.18)–(7.20) together with similar expressions for $\tilde{s}^2 \leftrightarrow x^2$ as for $\tilde{s}^1 \leftrightarrow x^1$.

Such equations provide a description of the conformal embedding which makes obvious the identification of antipodal points at infinity in $S_{[+],\kappa_2}^2$ because these are mapped by stereographical projection into the same image. Further, this description in ambient Weierstrass coordinates can be translated to any choice of intrinsic coordinate systems in either the initial space $S_{[\kappa_1],\kappa_2}^2$ or in the conformal compactified one $C_{[\kappa_1],\kappa_2}^2 \equiv \tilde{S}_{[+],\kappa_2}^2$. For the former, if in the 2D version of (7.20) we introduce the parametrizations given in table 2, we recover the results displayed in table 9. For the latter, considered as CK spaces in their natural coordinates, denoted here in capital letters (parallel I (A, Y) , parallel II (X, B) , polar (R, Φ) ; see table 2), the embedding into the ambient space $\mathbb{R}^3 = (\tilde{s}^+, \tilde{s}^1, \tilde{s}^2)$ is:

$$\begin{aligned} \tilde{s}^+ &= \cos A \, C_{\kappa_2}(Y) = \cos X \, C_{\kappa_2}(B) = \cos R \\ \tilde{s}^1 &= \sin A \, C_{\kappa_2}(Y) = \sin X = \sin R \, C_{\kappa_2}(\Phi) \\ \tilde{s}^2 &= S_{\kappa_2}(Y) = \cos X \, S_{\kappa_2}(B) = \sin R \, S_{\kappa_2}(\Phi). \end{aligned} \tag{7.28}$$

Thus we can find the description of the embedding $S_{[\kappa_1],\kappa_2}^2 \mapsto C_{[\kappa_1],\kappa_2}^2$ in any coordinates required that are also written in table 9. As an example, when parallel I type coordinates are used in both $S_{[\kappa_1],\kappa_2}^2$ and $\tilde{S}_{[+],\kappa_2}^2$, the explicit embeddings of the nine CK spaces are shown in table 10; note that whenever $\kappa_1\ell^2 = \{1, 0, -1\}$ the expressions in tables 9 and 10 are substantially simplified.

By taking into account the results given in table 10, the *conformal completion* for each CK space can clearly be visualized in the section $s^- = -1$ of the conformal ambient

space $\mathbb{R}^4 = (\tilde{s}^+, \tilde{s}^- = -1, \tilde{s}^1, \tilde{s}^2)$. The *conformal embedding* can be visualized through a stereographic projection with pole $\mathcal{P} = (-1, 0, 0)$ in the common space obtained by identifying the initial ambient CK space $(x^0, x^1/\ell, x^2/\ell)$ to the section $(\tilde{s}^+, \tilde{s}^1, \tilde{s}^2)$ of the conformal ambient space, as depicted for the 1D case in figure 2. In particular, the nine CK spaces are embedded according to the sign of κ_2 , into its three completions $\widetilde{\mathbf{S}^2} \equiv \mathbf{S}^2, \widetilde{\mathbf{NH}}_+^{1+1}, \widetilde{\mathbf{AdS}}^{1+1}$ as follows.

7.2.2 Riemannian spaces with $\kappa_2 > 0$: $S_{[\kappa_1],+}^2 \mapsto C_{[\kappa_1],+}^2 \equiv \mathbf{S}^2$

The embedding of the *sphere* $S_{[+],+}^2$ covers the full sphere $\mathbf{S}^2$ once, so that the sphere coincides with its conformal compactification. The embedding of the *Euclidean plane* $\mathbf{E}^2$ covers the full sphere $\mathbf{S}^2$ minus a single point, with conformal ambient coordinates $(-1, -1, 0, 0)$, that is, the pole $\mathcal{P}$ of the stereographic projection. This is the usual point at infinity required to globally define inversions and transforms $\mathbf{E}^2$ into the Riemann sphere. The embedding of the *Lobachewski plane* $\mathbf{H}^2$ only covers one half of the sphere ($\tilde{s}^+ > 0$ when ℓ is chosen such that $\kappa_1 \ell^2 = -1$), and points at infinity in $\mathbf{H}^2$ appear as ordinary points in the equator ($\tilde{s}^+ = 0$); the new points added in the compactification make up another half of the sphere ($\tilde{s}^+ < 0$), which may be considered as another copy of $\mathbf{H}^2$ glued to the proper embedding of the hyperbolic plane in $\mathbf{S}^2$ by the points in the equator $\tilde{s}^+ = 0$.

7.2.3 Non-relativistic spacetimes with $\kappa_2 = 0$: $S_{[\kappa_1],0}^2 \mapsto C_{[\kappa_1],0}^2 \equiv \widetilde{\mathbf{NH}}_+^{1+1}$

The transition from $\mathbf{NH}_+^{1+1}$ to $\widetilde{\mathbf{NH}}_+^{1+1}$ requires antipodal identification of the two circles $(\tilde{s}^+)^2 + (\tilde{s}^1)^2 = 1$ at the infinity $\tilde{s}^2 = \pm\infty$. Thus the new space is compact. Once this is assumed, the description of the compactification of the three non-relativistic spacetimes is straightforward. For *oscillating NH* spacetime $\mathbf{NH}_+^{1+1}$, one has simply to embed identically $\mathbf{NH}_+^{1+1}$ into itself. For the *Galilei* spacetime $\mathbf{G}^{1+1}$, its embedding into $\mathbf{NH}_+^{1+1}$ covers this space minus the line $\tilde{s}^+ = -1, \tilde{s}^- = -1, \tilde{s}^1 = 0$, any $\tilde{s}^2$; the (pole) point $\mathcal{P} = (-1, -1, 0, 0)$ on this line plays the role of point at infinity in $\mathbf{G}^{1+1}$ and other points on this line correspond to the ‘instantaneous 1D space’ of the point at infinity. The embedding of the *expanding NH* spacetime $\mathbf{NH}_-^{1+1}$ into $\mathbf{NH}_+^{1+1}$ covers one ‘upper’ half of the cylinder ($\tilde{s}^+ > 0$), so that one has to add as new points the remaining ‘lower’ half of the cylinder ($\tilde{s}^+ < 0$) glued to the other proper half by the two lines in the plane $\tilde{s}^+ = 0$. Further in the three cases one has to consider the antipodal identification of points at infinity in $\mathbf{NH}_+^{1+1}$ giving rise finally to $\widetilde{\mathbf{NH}}_+^{1+1}$ as the conformally compactified space.

7.2.4 Relativistic spacetimes with $\kappa_2 = -1/c^2 < 0$: $S_{[\kappa_1],-}^2 \mapsto C_{[\kappa_1],-}^2 \equiv \widetilde{\mathbf{AdS}}^{1+1}$

The transition from $\mathbf{AdS}^{1+1}$ to $\widetilde{\mathbf{AdS}}^{1+1}$ follows by antipodal identification of the two circles at the spatial infinity in the *anti-de Sitter* spacetime: $(\tilde{s}^+)^2 + (\tilde{s}^1)^2 - (\tilde{s}^2)^2/c^2 = 1$; hence $(\tilde{s}^+)^2 + (\tilde{s}^1)^2 = \infty, \tilde{s}^2 = \pm\infty$ and each circle $\tilde{s}^2 = \infty, \tilde{s}^2 = -\infty$ is antipodal to the other. The (compact) time-like lines (label $\kappa_1 > 0$) embed homeomorphically, while the space-like lines, whose label is $\kappa_1 \kappa_2 < 0$, hence hyperbolic and originally not compact, are glued by their points at infinity with their antipodals lines in $\mathbf{AdS}^{1+1}$, following

the antipodal identification of their points at infinity; hence they become also compact. Topologically the space so obtained is $\mathbf{S}^1 \times \mathbf{S}^1$. One $\mathbf{S}^1$ corresponds to the originally compact time-like line l_1 . The other comes from the 1D compactification of the originally hyperbolic, hence non-compact space-like line l_2 , which is obtained by glueing two copies of a hyperbolic line (l_2 and its antipodal) by their points at infinity, as discussed in the section 7.1.1.

For *Minkowskian* spacetime $\mathbf{M}^{1+1}$, the completion requires to add two lines with equations $\tilde{s}^+ = -1, \tilde{s}^- = -1, \tilde{s}^2 = \pm c \tilde{s}^1$ crossing through the intersection (pole) point $\mathcal{P}$. This is rather well known: the new point $\mathcal{P}$ corresponds to the point ‘at infinity’ in $\mathbf{M}^{1+1}$ and the two new lines are the light-cone of the point at infinity.

Finally, the *de Sitter* spacetime $\mathbf{dS}^{1+1}$, with hyperbolic non-compact time-like lines ($\kappa_1 < 0$) and compact space-like ones ($\kappa_1 \kappa_2 > 0$) is the only CK space where parallel I coordinates do not cover completely the space, so the expressions in table 10 do not provide a complete description. In this case it is better to use directly stereographic projection, which maps the whole de Sitter spacetime into $\mathbf{AdS}^{1+1}$ in a one-to-one way compatible with antipodal identification in $\mathbf{AdS}^{1+1}$. This could have been foreseen, as essentially $\mathbf{dS}^{1+1}$ and $\mathbf{AdS}^{1+1}$ are the same space, with an interchange time-like $\leftrightarrow$ space-like, so we can expect $\mathbf{dS}^{1+1}$ and $\mathbf{AdS}^{1+1}$ to have essentially the same compactification. If we are interested only in the double wedge in $\mathbf{dS}^{1+1}$ covered by parallel I coordinates (with focal points at the poles of the initial time-like line l_1), the embedding of this region is determined by $(\tilde{s}^1)^2 < 1$, that is, $0 < (\tilde{s}^+)^2 - (\tilde{s}^2)^2/c^2 \leq 1$, which is limited by four lines with equations $\tilde{s}^- = -1, \tilde{s}^1 = \pm 1, \tilde{s}^2 = \pm c \tilde{s}^+$; these four lines are the two pairs of isotropic lines through the two poles of the initial line l_1 . However, the other regions limited by these lines in the conformal compactification are not new points, but the images by the embedding of the region of $\mathbf{dS}^{1+1}$ not covered by the parallel I coordinates.

8 Concluding remarks

The present paper affords an approach to the conformal algebras, groups and spaces comprehensive enough to give a global understanding of those aspects of conformal invariance related with either the space curvature or the metric signature in the initial space. In particular, it produces a very explicit description of the *conformal completion* of the homogeneous spacetimes where the fact that the conformal completion of a homogeneous space of constant curvature κ_1 does not depend on the curvature can be clearly seen. This result may be foreseen from the well known fact that every 2D metric, constant curvature or not, is conformally flat—and hence *a fortiori* the conformal compactification of a curved space will coincide with that of its corresponding flat one—, but we have attempted here to make an *ab-initio* approach to cycle-preserving transformations because this provides a lot of additional details, and further can also be extended to higher dimensions, where situation is rather different. A topic has not deliberately been touched upon: the use of the three one-step Cayley–Dickson extensions of $\mathbb{R}$ (complex, dual or double numbers [13]), in terms of which circular transformations of the nine CK spaces can be represented as fractional linear (Möbius-like) transformations; this is because this possibility only exists for two dimensions.

In both Euclidean or Minkowski spaces the transition from the motion to the conformal group can be looked at in two stages: the motion group can be extended first by a

one-parameter dilation subgroup, obtaining a similitude group, and then by the specific conformal transformations, closing the whole conformal group. It is a widespread belief that dilation-like transformations do not exist in spaces with non-zero curvature (sphere, anti-de Sitter, ...), but the explicit results obtained here show this is not so. What is actually different for non-zero curvature is that the intermediate stage provided by similitudes does not exist, and once a single dilation is added to the motion group, the full conformal group is obtained. Another trait follows from the subalgebra structure. For $\kappa_1 = 0$ there is a complete symmetry between translation generators P_i and specific conformal generators G_i , and usually these are introduced as conjugate to translations by an inversion in the origin. This symmetry *does not* extend to the non-zero curvature case, and while the P_i *do not* commute among themselves, the G_i do always commute. There is however another *conformal duality* between translations and Λ -translations, with respective generators P_i , L_i , but this duality is invisible in the conformal algebra of flat spaces, where Λ -translations coincide with translations, leaving the specific conformal transformations as a kind of vestigial residue of the difference between P_i and L_i .

From a more general viewpoint, a study with κ_1, κ_2 as *parameters* displays many illuminating clues which are clear in this generic approach but cannot be seen in a single flat case. In previous papers [14, 16, 20, 21] we have established the suitability of this type of approach and have developed many of the geometrical and algebraic aspects involved. And further, as this approach has a built-in scheme of contractions, its separate study is made completely redundant.

Some problems in the conventional Minkowskian quantum field theories can be seen in a new light if dealt with in the non-zero curvature case (anti-de Sitter or de Sitter) taking afterwards a flat limit; in this sense to study dependence on the curvature is a natural inquiry. The structure of conformal compactification of Minkowskian spacetime appears as a particular instance within our parametric approach; it allows a clear understanding and visualization of how the embedding of the initial spacetime changes when its curvature vanishes or when the metric degenerates. The first issue is of relevance in view of the current interest in AdS-CFT correspondence as a conjecture relating local QFT on $\text{AdS}^{1+(d-1)}$ to a conformal QFT on the compactified Minkowski spacetime $\text{CompM}^{1+(d-2)}$ (see [33] and references therein). In this context an explicit description of the geometry behind these spaces in a way as general as possible should be helpful. In another context, the degeneration of a Lorentzian-type metric produces Newtonian theories, and ‘non-relativistic electromagnetic theories’, Maxwell–Le Bellac–Lévy-Leblond equations [26, 27], as non-relativistic limits of Maxwell equations also fit inside this parametrized scheme. Furthermore, the results here obtained constitute a starting point for the development of anti-de Sitter and de Sitter electromagnetic theories.

Finally, the techniques and results we have presented here can be generalized straightforwardly to higher dimensions, where the differential realizations of conformal groups can be obtained in a completely explicit and closed form for any curvature and signature. This will be summarized in a forthcoming paper.

Acknowledgments

This work was partially supported by the Ministerio de Ciencia y Tecnología, Spain (Projects PB98-0370 and BFM2000-1055) and by Junta de Castilla y León, Spain (Project CO2/399).

Appendix A. The Lambda function

Let us recall the definitions for the Gudermannian function $\text{gd}(x)$ and the function $\lambda(x)$ [34] which often appear in hyperbolic geometry:

$$\text{gd}(x) = \frac{\pi}{2} - 2 \arctan(e^{-x}) = 2 \arctan(e^x) - \frac{\pi}{2} \quad (\text{A.1})$$

$$\lambda(x) = -i \frac{\pi}{2} - 2 \operatorname{arctanh}(-ie^{-ix}) = 2i \arctan(e^{-ix}) - i \frac{\pi}{2}. \quad (\text{A.2})$$

They are related by $\lambda(x) = i \text{gd}(-ix)$ and they are inverse of each other, $\text{gd}(\lambda(x)) = x, \lambda(\text{gd}(x)) = x$. Alternatively, these functions may be defined by the functional relations:

$$\tanh\left(\frac{\lambda(x)}{2}\right) = \tan\left(\frac{x}{2}\right) \quad \tanh\left(\frac{x}{2}\right) = \tan\left(\frac{\text{gd}(x)}{2}\right) \quad (\text{A.3})$$

showing that if $x \in (-\infty, \infty)$, then $\text{gd}(x) \in (-\pi/2, \pi/2)$. If $\text{gd}(x)$ is considered as a point in the circle $\mathbf{S}^1 \equiv (-\pi, \pi]$, the image of $\mathbb{R}$ by the map $\text{gd}(x)$ only fills half the circle. Alternatively, the map $\lambda(x)$ is only defined in half the circle $x \in (-\pi/2, \pi/2)$, and the image is the whole line $\mathbb{R}$.

Within the parametrized CK approach it is natural to consider the *Lambda function* $\Lambda_\kappa(x)$ defined as:

$$\Lambda_\kappa(x) = \int_0^x \frac{1}{C_\kappa(t)} dt. \quad (\text{A.4})$$

By taking into account the derivative of $\operatorname{arcT}_\kappa(x)$ (3.5) we find:

$$\int_0^x \frac{1}{C_\kappa(t)} dt = \left[-2 \operatorname{arcT}_{-\kappa} \left(\frac{1}{\sqrt{-\kappa}} e^{-\sqrt{-\kappa}t} \right) \right]_{t=0}^{t=x} \quad (\text{A.5})$$

and since

$$\operatorname{arcT}_{-\kappa}(1/\sqrt{-\kappa}) = \frac{1}{\sqrt{-\kappa}} \frac{\pi}{4} \quad (\text{A.6})$$

we obtain

$$\Lambda_\kappa(x) = \frac{1}{\sqrt{-\kappa}} \frac{\pi}{2} - 2 \operatorname{arcT}_{-\kappa} \left(\frac{1}{\sqrt{-\kappa}} e^{-\sqrt{-\kappa}x} \right) = \begin{cases} \frac{1}{\sqrt{\kappa}} \lambda(\sqrt{\kappa}x) & \kappa > 0 \\ x & \kappa = 0 \\ \frac{1}{\sqrt{-\kappa}} \text{gd}(\sqrt{-\kappa}x) & \kappa < 0 \end{cases} \quad (\text{A.7})$$

Therefore both the lambda function (A.2) and the gudermannian (A.1) are the two particular elliptic $\kappa > 0$ or hyperbolic $\kappa < 0$ instances of a single CK labelled ‘Lambda function’: $\Lambda_{+1}(x) \equiv \lambda(x)$ and $\Lambda_{-1}(x) \equiv \text{gd}(x)$. The analogous of the functional definition (A.3), which can be obtained starting from (A.7), reads:

$$T_{-\kappa} \left(\frac{\Lambda_\kappa(x)}{2} \right) = T_\kappa \left(\frac{x}{2} \right) \quad T_{-\kappa} \left(\frac{x}{2} \right) = T_\kappa \left(\frac{\Lambda_{-\kappa}(x)}{2} \right) \quad (\text{A.8})$$

whence

$$\Lambda_{-\kappa}(\Lambda_\kappa(x)) = x. \quad (\text{A.9})$$

This property extends the known fact that $\lambda(x)$ and $\text{gd}(x)$ are inverse each other.

From this viewpoint, if x is a quantity with label κ , i.e. a point in the 1D CK space $S_{[\kappa]}^1$, then $\Lambda_\kappa(x)$ (also denoted $x^\wedge$) is a quantity with label $-\kappa$, i.e. a point in $S_{[-\kappa]}^1$, and

the Lambda function should be seen as providing a canonical identification between $S_{[\kappa]}^1$ and $S_{[-\kappa]}^1$, that is, between quantities with elliptic and hyperbolic labels. In particular, from (A.8) it can be found that

$$C_{-\kappa}(\Lambda_\kappa(x)) = \frac{1}{C_\kappa(x)} \quad S_{-\kappa}(\Lambda_\kappa(x)) = T_\kappa(x) \quad T_{-\kappa}(\Lambda_\kappa(x)) = S_\kappa(x) \quad (\text{A.10})$$

$$C_{-\kappa}^2(\Lambda_\kappa(x)) - \kappa S_{-\kappa}^2(\Lambda_\kappa(x)) = 1 \quad (\text{A.11})$$

and thus $\Lambda_\kappa(x)$ has label $-\kappa$. The derivative of $\Lambda_\kappa(x)$ is:

$$\frac{d\Lambda_\kappa(x)}{dx} = \frac{1}{C_\kappa(x)} = C_{-\kappa}(\Lambda_\kappa(x)). \quad (\text{A.12})$$

As commented before, for *positive* κ the Lambda function is *not* defined in the whole CK space $S_{[\kappa]}^1$. However this function can be extended in a natural way to a function, still denoted under the same name, which is defined in the whole $S_{[\kappa]}^1$. In this way the image of the elliptic 1D space $S_{[\kappa]}^1$ (with $\kappa > 0$) by Λ_κ should be identified with *two* copies of the hyperbolic 1D space $S_{[-\kappa]}^1$ which are glued by their final points. The transcription of the geometrical approach done in the paper to an algebraic approach involves the use of this extension of the Lambda function.

The Lambda function has a clear geometrical interpretation in the anti-de Sitter, Minkowskian and de Sitter spacetimes with curvature κ_1 and $\kappa_2 = -1/c^2$: if t (label κ_1) is a time-like length, and we build an orthogonal triangle with an isotropic hypotenuse (angle $\equiv$ rapidity equals infinity), then the space length of the other side is $c\Lambda_{\kappa_1}(t)$ [15]. In particular, in Minkowskian spacetime, $\kappa_1 = 0$, this reduces to the well known ct .

Appendix B. Geodesic curvature and finite distance

In a 2D space, the geodesic curvature of a curve can be defined as $k_g = |\frac{d\beta}{ds}|$, where $\beta(s)$ is the angle between any vector in parallel transport along the line and the tangent vector to the curve at the point with canonical parameter s [10]. In terms of the metric of a 2D space $ds^2 = \sum_{i,j=1}^2 g_{ij} du^i du^j$ and connexion symbols Γ_{jk}^i , the geodesic curvature for a line $(u^1(s), u^2(s))$ is known to be

$$k_g^2 = \sum_{i,j=1}^2 g_{ij} v^i v^j \quad v^i = \frac{d^2 u^i}{ds^2} + \sum_{j,k=1}^2 \Gamma_{jk}^i \frac{du^j}{ds} \frac{du^k}{ds} \quad i = 1, 2. \quad (\text{B.1})$$

We proceed to deduce the general expression of k_g for a cycle in parallel I coordinates $(u^1, u^2) = (a, y)$. By taking into account the cycle equation (4.2) and the main metric $(ds)_1^2$ written in table 2, and using the shorthand

$$f(a, y) = \frac{-\alpha_0 \kappa_1 S_{\kappa_1}(a) + \alpha_1 C_{\kappa_1}(a)}{\alpha \kappa_1 \kappa_2 S_{\kappa_1 \kappa_2}(y) - \alpha_2} C_{\kappa_1 \kappa_2}(y)$$

we find for the first derivatives

$$\frac{da}{ds} = \frac{1}{C_{\kappa_1 \kappa_2}(y)} (1 + \kappa_2 f^2(a, y))^{-1/2} \quad \frac{dy}{ds} = f(a, y) (1 + \kappa_2 f^2(a, y))^{-1/2}$$

(implying $\frac{dy}{da} = f(a, y)C_{\kappa_1\kappa_2}(y)$) and for the second derivatives

$$\begin{aligned}\frac{d^2a}{ds^2} &= \kappa_1\kappa_2 \frac{f(a, y)}{C_{\kappa_1\kappa_2}^2(y)} \left(S_{\kappa_1\kappa_2}(y) + \frac{\alpha - \alpha_2 S_{\kappa_1\kappa_2}(y)}{\alpha\kappa_1\kappa_2 S_{\kappa_1\kappa_2}(y) - \alpha_2} \right) (1 + \kappa_2 f^2(a, y))^{-1} \\ \frac{d^2y}{ds^2} &= -\frac{\kappa_1}{C_{\kappa_1\kappa_2}(y)} \left(\frac{\alpha - \alpha_2 S_{\kappa_1\kappa_2}(y)}{\alpha\kappa_1\kappa_2 S_{\kappa_1\kappa_2}(y) - \alpha_2} \right) (1 + \kappa_2 f^2(a, y))^{-1}.\end{aligned}$$

Therefore, in this case the two components v^i (B.1) turn out to be

$$\begin{aligned}v^1 &= \frac{d^2a}{ds^2} - 2\kappa_1\kappa_2 T_{\kappa_1\kappa_2}(y) \frac{da}{ds} \frac{dy}{ds} = \frac{\alpha\kappa_1\kappa_2 f(a, y)}{\alpha\kappa_1\kappa_2 S_{\kappa_1\kappa_2}(y) - \alpha_2} (1 + \kappa_2 f^2(a, y))^{-1} \\ v^2 &= \frac{d^2y}{ds^2} + \kappa_1 S_{\kappa_1\kappa_2}(y) C_{\kappa_1\kappa_2}(y) \left(\frac{da}{ds} \right)^2 = -\frac{\alpha\kappa_1 C_{\kappa_1\kappa_2}(y)}{\alpha\kappa_1\kappa_2 S_{\kappa_1\kappa_2}(y) - \alpha_2} (1 + \kappa_2 f^2(a, y))^{-1}\end{aligned}$$

so k_g is finally obtained as the length of the vector v^i in the main metric:

$$k_g^2 = C_{\kappa_1\kappa_2}^2(y)(v^1)^2 + \kappa_2(v^2)^2 = \frac{\kappa_1^2\kappa_2\alpha^2}{\alpha_2^2 + \kappa_2\alpha_1^2 + \kappa_1\kappa_2\alpha_0^2 - \kappa_1\kappa_2\alpha^2} \quad (\text{B.2})$$

where we have used the cycle equation and the identity

$$\kappa_1(\alpha_0 C_{\kappa_1}(a) + \alpha_1 S_{\kappa_1}(a))^2 + (-\alpha_0 \kappa_1 S_{\kappa_1}(a) + \alpha_1 C_{\kappa_1}(a))^2 = \alpha_1^2 + \kappa_1 \alpha_0^2. \quad (\text{B.3})$$

If $\alpha = 0$, then $v^i = 0$ reproduces the differential equations of geodesics with $k_g = 0$.

On the other hand, the finite form of the ('time-like') distances associated to the main metric $(ds)_1$ between two points (given in table 5) can be obtained by integrating $(ds)_1$ along the geodesic through the points. Working again with parallel I coordinates and considering the corresponding geodesic equation in table 5 and the identity (B.3), one finds:

$$ds = \frac{(1 + \kappa_1\kappa_2\beta_0^2 + \kappa_2\beta_1^2)^{1/2}}{1 + \kappa_1\kappa_2(\beta_0 C_{\kappa_1}(a) + \beta_1 S_{\kappa_1}(a))^2} da.$$

Hence the distance between two points with parallel coordinates (a_0, y_0) and (a_1, y_1) reads

$$s = \left[\text{arcT}_{\kappa_1} \left(\frac{\kappa_2\beta_0\beta_1 + (1 + \kappa_2\beta_1^2)T_{\kappa_1}(a)}{(1 + \kappa_1\kappa_2\beta_0^2 + \kappa_2\beta_1^2)^{1/2}} \right) \right]_{a=a_0}^{a=a_1}.$$

Since both points must lie on the same geodesic, $T_{\kappa_1\kappa_2}(y_i) = \beta_0 C_{\kappa_1}(a_i) + \beta_1 S_{\kappa_1}(a_i)$ ($i = 0, 1$), we may obtain the constants β_0, β_1 in terms of the end points as:

$$\beta_0 = \frac{S_{\kappa_1}(a_1)T_{\kappa_1\kappa_2}(y_0) - S_{\kappa_1}(a_0)T_{\kappa_1\kappa_2}(y_1)}{S_{\kappa_1}(a_1 - a_0)} \quad \beta_1 = \frac{C_{\kappa_1}(a_0)T_{\kappa_1\kappa_2}(y_1) - C_{\kappa_1}(a_1)T_{\kappa_1\kappa_2}(y_0)}{S_{\kappa_1}(a_1 - a_0)}.$$

Therefore the expression for the tangent of the distance is given by

$$T_{\kappa_1}(s) = \frac{S_{\kappa_1}(a_1 - a_0)(1 + \kappa_1\kappa_2\beta_0^2 + \kappa_2\beta_1^2)^{1/2}}{C_{\kappa_1}(a_1 - a_0) + \kappa_1\kappa_2 T_{\kappa_1\kappa_2}(y_0)T_{\kappa_1\kappa_2}(y_1)}$$

and the expression for the cosine of the distance given in table 5 is directly recovered

$$C_{\kappa_1}^2(s) = \frac{1}{1 + \kappa_1 T_{\kappa_1}^2(s)} = (C_{\kappa_1\kappa_2}(y_0)C_{\kappa_1\kappa_2}(y_1)C_{\kappa_1}(a_1 - a_0) + \kappa_1\kappa_2 S_{\kappa_1\kappa_2}(y_0)S_{\kappa_1\kappa_2}(y_1))^2. \quad (\text{B.4})$$

An alternative trigonometric procedure to find the finite ('time-like') distance between two points in polar coordinates (r_0, ϕ_0) and (r_1, ϕ_1) is to consider the cosine theorem for a side c in a CK triangle as is given in the equations (4.4) of [16]:

$$C_{\kappa_1}(c) = C_{\kappa_1}(a)C_{\kappa_1}(b) + \kappa_1 S_{\kappa_1}(a)S_{\kappa_1}(b)C_{\kappa_2}(C).$$

Taking the vertex C as the origin, then the sides a and b are the coordinates r_0 and r_1 , the angle C is the difference $\phi_1 - \phi_0$ and the side c is the distance s between both points:

$$C_{\kappa_1}(s) = C_{\kappa_1}(r_0)C_{\kappa_1}(r_1) + \kappa_1 S_{\kappa_1}(r_0)S_{\kappa_1}(r_1)C_{\kappa_2}(\phi_1 - \phi_0). \quad (\text{B.5})$$

This is the expression given in table 5 and it is equivalent to (B.4).

Appendix C. Conformal transformations of the metrics

To obtain the vector fields X corresponding to the conformal generators of a given metric g , the usual method is to enforce the proportionality by a positive factor $\mu(x)$ between g and the Lie derivative of g ; this leads to the *conformal Killing equations* of the metric (for the flat spaces $ISO(p, q)/SO(p, q)$ this approach can be found in [10]). We study here such transformations for the metrics g_1 and g_2 in the spaces $S_{[\kappa_1], \kappa_2}^2$. Let g be the metric of a 2D space with coordinates (u^1, u^2) : $ds^2 = \sum_{i,j=1}^2 g_{ij} du^i du^j$. The Lie derivative $L_X g$ of g , when X is a generator of a Lie group G acting on the space is [35]:

$$L_X g = \sum_{i,j,k=1}^2 \left(g_{ik} \partial_j X^k + g_{kj} \partial_i X^k + X^k \partial_k g_{ij} \right) du^i du^j \quad (\text{C.1})$$

where X^k are the components of the generator X written as a vector field: $X = \sum_k X^k \partial_k$.

Let us consider first the main metric g_1 of the spaces $S_{[\kappa_1], \kappa_2}^2$ in parallel I coordinates as given in table 2. The non-zero components of g_1 are $g_{aa} = C_{\kappa_1 \kappa_2}^2(y)$ and $g_{yy} = \kappa_2$, so that for a generic vector field $X \equiv X^a \partial_a + X^y \partial_y$ the Lie derivative (C.1) reads:

$$\begin{aligned} L_X g_1 = & 2 \left(C_{\kappa_1 \kappa_2}^2(y) \partial_a X^a - \kappa_1 \kappa_2 X^y C_{\kappa_1 \kappa_2}(y) S_{\kappa_1 \kappa_2}(y) \right) da^2 \\ & + 2 \kappa_2 (\partial_y X^y) dy^2 + 2 \left(C_{\kappa_1 \kappa_2}^2(y) \partial_y X^a + \kappa_2 \partial_a X^y \right) da dy. \end{aligned} \quad (\text{C.2})$$

If X is the vector field of the generators $\{P_1, P_2, J_{12}\}$ given in table 7, the expression (C.2) vanishes, in plain concordance with their role as isometries of $S_{[\kappa_1], \kappa_2}^2$. Let us see what happen with the three remaining generators of $\text{conf}_{\kappa_1, \kappa_2}$. The components of the dilation generator D are

$$D^a = -\frac{S_{\kappa_1}(a)}{C_{\kappa_1 \kappa_2}(y)} \quad D^y = -C_{\kappa_1}(a) S_{\kappa_1 \kappa_2}(y). \quad (\text{C.3})$$

By taking into account (3.5) we obtain

$$\begin{aligned} \partial_a D^a &= -\frac{C_{\kappa_1}(a)}{C_{\kappa_1 \kappa_2}(y)} & \partial_y D^a &= -\kappa_1 \kappa_2 \frac{S_{\kappa_1}(a) S_{\kappa_1 \kappa_2}(y)}{C_{\kappa_1 \kappa_2}^2(y)} \\ \partial_a D^y &= \kappa_1 S_{\kappa_1}(a) S_{\kappa_1 \kappa_2}(y) & \partial_y D^y &= -C_{\kappa_1}(a) C_{\kappa_1 \kappa_2}(y) \end{aligned} \quad (\text{C.4})$$

and by substituting this result in (C.2) we finally have

$$L_D g_1 = -2 C_{\kappa_1}(a) C_{\kappa_1 \kappa_2}(y) \left(C_{\kappa_1 \kappa_2}^2(y) da^2 + \kappa_2 dy^2 \right) = -2 C_{\kappa_1}(a) C_{\kappa_1 \kappa_2}(y) g_1. \quad (\text{C.5})$$

Therefore, the Lie derivative of the main metric along the dilation generator is proportional to g_1 by a *conformal factor*:

$$\mu_D(a, y) = -2C_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y). \quad (\text{C.6})$$

Likewise, we find the conformal factors associated to the generators G_i :

$$\mu_{G_1}(a, y) = 2S_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y) \quad \mu_{G_2}(a, y) = 2\kappa_2 S_{\kappa_1\kappa_2}(y). \quad (\text{C.7})$$

For the the subsidiary metric, we recall that g_2 is only relevant whenever $\kappa_2 = 0$, and in this case it is a metric defined only in each foliation leaf $a = \text{constant}$, with expression $(ds^2)_2 = dy^2$. It is straightforward to check that the results obtained starting from g_2 fully coincide with those obtained for g_1 by means of the contraction $\kappa_2 \rightarrow 0$: $\mu_{P_1} = \mu_{P_2} = \mu_{J_{12}} = \mu_{G_2} = 0$, $\mu_D = -2C_{\kappa_1}(a)$ and $\mu_{G_1} = 2S_{\kappa_1}(a)$; this could also have been derived from the fact that g_2 is generically proportional to g_1 .

Hence we conclude that both metrics verify

$$L_X g_i = \mu_X(u^1, u^2) g_i \quad X \in \{P_i, J_{12}, G_i, D\} \quad i = 1, 2 \quad (\text{C.8})$$

where the conformal factors $\mu_X(u^1, u^2)$ are given in terms of the three intrinsic geodesic and Weierstrass coordinates by

$$\begin{aligned} \mu_{P_1} &= \mu_{P_2} = \mu_{J_{12}} = 0 \\ \mu_D &= -2C_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y) = -2C_{\kappa_1}(x)C_{\kappa_1\kappa_2}(b) = -2C_{\kappa_1}(r) \equiv -2x^0 \\ \mu_{G_1} &= 2S_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y) = 2S_{\kappa_1}(x) = 2S_{\kappa_1}(r)C_{\kappa_2}(\phi) \equiv 2x^1 \\ \mu_{G_2} &= 2\kappa_2 S_{\kappa_1\kappa_2}(y) = 2\kappa_2 C_{\kappa_1}(x)S_{\kappa_1\kappa_2}(b) = 2\kappa_2 S_{\kappa_1}(r)S_{\kappa_2}(\phi) \equiv 2\kappa_2 x^2. \end{aligned} \quad (\text{C.9})$$

When $\kappa_1 = 0$ we recover the known results concerning $\mathbf{E}^2$ and $\mathbf{M}^{1+1}$ [10]:

$$\begin{aligned} \mu_{P_1} &= \mu_{P_2} = \mu_{J_{12}} = 0 & \mu_{G_1} &= 2a = 2x = 2rC_{\kappa_2}(\phi) \\ \mu_D &= -2 & \mu_{G_2} &= 2\kappa_2 y = 2\kappa_2 b = 2\kappa_2 rS_{\kappa_2}(\phi). \end{aligned} \quad (\text{C.10})$$

Notice that the conformal factors (C.9) coincide with the factors multiplying $\mathcal{C}$ in (6.4); this could be expected since the metrics in the CK space are related with the second-order Casimir $\mathcal{C}$ through the Killing–Cartan form.

References

- [1] Cunningham E 1910 *Proc. London Math. Soc.* **8** 77
- [2] Bateman H 1910 *Proc. London Math. Soc.* **8** 223
- [3] Rosen G 1972 *Am. J. Phys.* **40** 1023
- [4] Fushchich W I and Nikitin A G 1987 *Symmetries of Maxwell's Equations* (Dordrecht: Reidel)
- [5] McLennan J A 1956 *Nuovo Cimento* **3** 1360
- [6] McLennan J A 1957 *Nuovo Cimento* **5** 640

- [7] Mack G and Salam A 1969 *Ann. Phys.* **53** 174
- [8] Mickelsson J and Niederle J 1975 *Ann. Inst. H. Poincaré* **23** 277
- [9] Jacobsen H P and Vergne M 1977 *J. Funct. Anal.* **24** 52
- [10] Doubrovine B, Novikov S and Fomenko A 1982 *Géométrie Contemporaine, Méthodes et Applications* First Part (Moscow: MIR)
- [11] Berger M 1987 *Geometry I* (Berlin: Springer)
- [12] Yaglom I M, Rozenfel'd B A and Yasinskaya E U 1966 *Sov. Math. Surveys* **19** 49
- [13] Yaglom I M 1979 *A Simple Non-Euclidean Geometry and its Physical Basis* (New York: Springer)
- [14] Ballesteros A, Herranz F J, del Olmo M A and Santander M 1993 *J. Phys. A: Math. Gen.* **26** 5801
- [15] Herranz F J 1995 Grupos de Cayley–Klein clásicos y cuánticos *PhD Thesis* University of Valladolid
- [16] Herranz F J, Ortega R and Santander M 2000 *J. Phys. A: Math. Gen.* **33** 4525
- [17] Rozenfel'd B A 1988 *A History of Non-Euclidean Geometry* (New York: Springer)
- [18] Rozenfel'd B A 1997 *Geometry of Lie groups* (Dordrecht: Kluwer)
- [19] Herranz F J and Santander M 1997 *J. Phys. A: Math. Gen.* **30** 5411
- [20] Ballesteros A, Herranz F J, del Olmo M A and Santander M 1995 *J. Math. Phys.* **36** 631
- [21] Rañada M F and Santander M 1999 *J. Math. Phys.* **40** 5026
- [22] Bacry H and Lévy-Leblond J M 1968 *J. Math. Phys.* **9** 1605
- [23] Herranz F J and Santander M 1998 Homogeneous phase spaces: the Cayley–Klein approach *Geometry and Physics* vol XXXII, ed J F Cariñena *et al* (Madrid: Real Academia de Ciencias) pp 59
- [24] Kobayashi S and Nomizu K 1963 *Foundations of Differential Geometry* Vol 1 (New York: Wiley-Interscience)
- [25] Klingenberg W 1978 *A Course in Differential Geometry* Graduate texts in Mathematics (New York: Springer)
- [26] Lévy-Leblond J M 1967 *Comm. Math. Phys.* **6** 286
- [27] Le Bellac M and Lévy-Leblond J M 1973 *Il Nuovo Cimento* **14B**(2) 217
- [28] Negro J, del Olmo M A and Rodríguez-Marco A 1997 *J. Math. Phys.* **38** 3786
- [29] Negro J, del Olmo M A and Rodríguez-Marco A 1997 *J. Math. Phys.* **38** 3810
- [30] Barut A O and Xu B 1981 *Phys. Lett. A* **82** 218

- [31] de Alfaro V, Fubini S, Furlan G 1978 Conformal invariance in field theory *Lect. Notes Math.* **676** 255 (1977 Differ. Geom. Meth. Math. Phys. II, Proc. Bonn)
- [32] Herranz F J and Santander M 1994 Conformal geometries for simple and quasisimple orthogonal groups *Differential Geometry and its Applications* (Anales de Física. Monografías 2) ed M A Cañadas-Pinedo *et al* (Madrid: CIEMAT-RSEF) pp 17
- [33] Rehren K H 2000 A proof of the AdS-CFT Correspondence *Quantum Theory and Symmetries* ed H D Doebner *et al* (Singapore: World Scientific) pp 278
- [34] Dwight H B 1969 *Tables of integrals and other mathematical data* (London: MacMillan) pp 150
- [35] Warner F W 1971 *Foundations of Differentiable Manifolds and Lie Groups* (Scott, Foresman and Company)

Table 1: The nine two-dimensional CK spaces $S^2_{[\kappa_1], \kappa_2} = SO_{\kappa_1, \kappa_2}(3)/SO_{\kappa_2}(2)$.

Elliptic: $\mathbf{S}^2$ $S^2_{[+],+} = SO(3)/SO(2)$	Euclidean: $\mathbf{E}^2$ $S^2_{[0],+} = ISO(2)/SO(2)$	Hyperbolic: $\mathbf{H}^2$ $S^2_{[-],+} = SO(2,1)/SO(2)$
Oscillating NH: $\mathbf{NH}^{1+1}_+$ (Co-Euclidean) $S^2_{[+],0} = ISO(2)/ISO(1)$	Galilean: $\mathbf{G}^{1+1}$ $S^2_{[0],0} = IISO(1)/ISO(1)$	Expanding NH: $\mathbf{NH}^{1+1}_-$ (Co-Minkowskian) $S^2_{[-],0} = ISO(1,1)/ISO(1)$
Anti-de Sitter: $\mathbf{AdS}^{1+1}$ (Co-Hyperbolic) $S^2_{[+],-} = SO(2,1)/SO(1,1)$	Minkowskian: $\mathbf{M}^{1+1}$ $S^2_{[0],-} = ISO(1,1)/SO(1,1)$	De Sitter: $\mathbf{dS}^{1+1}$ (Doubly Hyperbolic) $S^2_{[-],-} = SO(2,1)/SO(1,1)$

Table 2: Parametrized expressions for Weierstrass coordinates, metric, canonical connection and area element for $S^2_{[\kappa_1], \kappa_2}$ given in the three systems of geodesic coordinates. The subsidiary metric $(ds^2)_2$ is only relevant whenever $\kappa_2 = 0$.

Parallel I (a, y)	Parallel II (x, b)	Polar (r, ϕ)
$x^0 = C_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y)$	$x^0 = C_{\kappa_1}(x)C_{\kappa_1\kappa_2}(b)$	$x^0 = C_{\kappa_1}(r)$
$x^1 = S_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y)$	$x^1 = S_{\kappa_1}(x)$	$x^1 = S_{\kappa_1}(r)C_{\kappa_2}(\phi)$
$x^2 = S_{\kappa_1\kappa_2}(y)$	$x^2 = C_{\kappa_1}(x)S_{\kappa_1\kappa_2}(b)$	$x^2 = S_{\kappa_1}(r)S_{\kappa_2}(\phi)$
$(ds^2)_1 = C_{\kappa_1\kappa_2}^2(y)da^2 + \kappa_2 dy^2$	$(ds^2)_1 = dx^2 + \kappa_2 C_{\kappa_1}^2(x)db^2$	$(ds^2)_1 = dr^2 + \kappa_2 S_{\kappa_1}^2(r)d\phi^2$
$(ds^2)_2 = dy^2$ for $a = a_0$	$(ds^2)_2 = C_{\kappa_1}^2(x)db^2$ for $x = x_0$	$(ds^2)_2 = S_{\kappa_1}^2(r)d\phi^2$ for $r = r_0$
$\Gamma_{aa}^y = \kappa_1 S_{\kappa_1\kappa_2}(y)C_{\kappa_1\kappa_2}(y)$	$\Gamma_{bb}^x = \kappa_1 \kappa_2 S_{\kappa_1}(x)C_{\kappa_1}(x)$	$\Gamma_{\phi\phi}^r = -\kappa_2 S_{\kappa_1}(r)C_{\kappa_1}(r)$
$\Gamma_{ay}^a = -\kappa_1 \kappa_2 T_{\kappa_1\kappa_2}(y)$	$\Gamma_{bx}^b = -\kappa_1 T_{\kappa_1}(x)$	$\Gamma_{\phi r}^\phi = 1/T_{\kappa_1}(r)$
$dS = C_{\kappa_1\kappa_2}(y) da \wedge dy$	$dS = C_{\kappa_1}(x) dx \wedge db$	$dS = S_{\kappa_1}(r) dr \wedge d\phi$

Table 3: Particularized expressions for Weierstrass coordinates, metric, canonical connection and area element for the nine spaces $S^2_{[\kappa_1], \kappa_2}$ in geodesic parallel I coordinates (a, y) . In the three Riemannian spaces (upper line) κ_1, κ_2 are normalized to $\kappa_1 \in \{1, 0, -1\}$ and $\kappa_2 = 1$. In the six spacetimes, $\kappa_1 = \pm 1/\tau^2$, $\kappa_2 = -1/c^2$, $a \equiv t$ is the time coordinate and y is the space one.

$\mathbf{S}^2 = S^2_{[+],+}$ $x^0 = \cos a \cos y$ $x^1 = \sin a \cos y$ $x^2 = \sin y$ $(ds^2)_1 = \cos^2 y da^2 + dy^2$ $\Gamma^y_{aa} = \sin y \cos y$ $\Gamma^a_{ay} = -\tan y$ $d\mathcal{S} = \cos y da \wedge dy$	$\mathbf{E}^2 = S^2_{[0],+}$ $x^0 = 1$ $x^1 = a$ $x^2 = y$ $(ds^2)_1 = da^2 + dy^2$ $\Gamma^y_{aa} = 0$ $\Gamma^a_{ay} = 0$ $d\mathcal{S} = da \wedge dy$	$\mathbf{H}^2 = S^2_{[-],+}$ $x^0 = \cosh a \cosh y$ $x^1 = \sinh a \cosh y$ $x^2 = \sinh y$ $(ds^2)_1 = \cosh^2 y da^2 + dy^2$ $\Gamma^y_{aa} = -\sinh y \cosh y$ $\Gamma^a_{ay} = \tanh y$ $d\mathcal{S} = \cosh y da \wedge dy$
$\mathbf{NH}^{1+1}_+ = S^2_{[+1/\tau^2],0}$ $x^0 = \cos(t/\tau)$ $x^1 = \tau \sin(t/\tau)$ $x^2 = y$ $(ds^2)_1 = dt^2$ $(ds^2)_2 = dy^2 \quad t = t_0$ $\Gamma^y_{tt} = \frac{1}{\tau^2} y \quad \Gamma^t_{ty} = 0$ $d\mathcal{S} = dt \wedge dy$	$\mathbf{G}^{1+1} = S^2_{[0],0}$ $x^0 = 1$ $x^1 = t$ $x^2 = y$ $(ds^2)_1 = dt^2$ $(ds^2)_2 = dy^2 \quad t = t_0$ $\Gamma^y_{tt} = 0 \quad \Gamma^t_{ty} = 0$ $d\mathcal{S} = dt \wedge dy$	$\mathbf{NH}^{1+1}_- = S^2_{[-1/\tau^2],0}$ $x^0 = \cosh(t/\tau)$ $x^1 = \tau \sinh(t/\tau)$ $x^2 = y$ $(ds^2)_1 = dt^2$ $(ds^2)_2 = dy^2 \quad t = t_0$ $\Gamma^y_{tt} = -\frac{1}{\tau^2} y \quad \Gamma^t_{ty} = 0$ $d\mathcal{S} = dt \wedge dy$
$\mathbf{AdS}^{1+1} = S^2_{[+1/\tau^2], -1/c^2}$ $x^0 = \cos(t/\tau) \cosh(y/c\tau)$ $x^1 = \tau \sin(t/\tau) \cosh(y/c\tau)$ $x^2 = c\tau \sinh(y/c\tau)$ $(ds^2)_1 = \cosh^2(y/c\tau) dt^2 - \frac{1}{c^2} dy^2$ $\Gamma^y_{tt} = \frac{c}{\tau} \sinh(y/c\tau) \cosh(y/c\tau)$ $\Gamma^t_{ty} = \frac{1}{c\tau} \tanh(y/c\tau)$ $d\mathcal{S} = \cosh(y/c\tau) dt \wedge dy$	$\mathbf{M}^{1+1} = S^2_{[0], -1/c^2}$ $x^0 = 1$ $x^1 = t$ $x^2 = y$ $(ds^2)_1 = dt^2 - \frac{1}{c^2} dy^2$ $\Gamma^y_{tt} = 0$ $\Gamma^t_{ty} = 0$ $d\mathcal{S} = dt \wedge dy$	$\mathbf{dS}^{1+1} = S^2_{[-1/\tau^2], -1/c^2}$ $x^0 = \cosh(t/\tau) \cos(y/c\tau)$ $x^1 = \tau \sinh(t/\tau) \cos(y/c\tau)$ $x^2 = c\tau \sin(y/c\tau)$ $(ds^2)_1 = \cos^2(y/c\tau) dt^2 - \frac{1}{c^2} dy^2$ $\Gamma^y_{tt} = -\frac{c}{\tau} \sin(y/c\tau) \cos(y/c\tau)$ $\Gamma^t_{ty} = -\frac{1}{c\tau} \tan(y/c\tau)$ $d\mathcal{S} = \cos(y/c\tau) dt \wedge dy$

Table 4: Same information as in table 3, with the same conventions, for the nine spaces $S^2_{[\kappa_1], \kappa_2}$ in geodesic polar coordinates (r, ϕ) . In the six spacetimes, r is a time-like length (a proper time), while $\phi \equiv \chi$ is a rapidity.

$\mathbf{S}^2 = S^2_{[+], +}$ $x^0 = \cos r$ $x^1 = \sin r \cos \phi$ $x^2 = \sin r \sin \phi$ $(ds^2)_1 = dr^2 + \sin^2 r d\phi^2$ $\Gamma^r_{\phi\phi} = -\sin r \cos r$ $\Gamma^\phi_{\phi r} = 1/\tan r$ $dS = \sin r dr \wedge d\phi$	$\mathbf{E}^2 = S^2_{[0], +}$ $x^0 = 1$ $x^1 = r \cos \phi$ $x^2 = r \sin \phi$ $(ds^2)_1 = dr^2 + r^2 d\phi^2$ $\Gamma^r_{\phi\phi} = -r$ $\Gamma^\phi_{\phi r} = 1/r$ $dS = r dr \wedge d\phi$	$\mathbf{H}^2 = S^2_{[-], +}$ $x^0 = \cosh r$ $x^1 = \sinh r \cos \phi$ $x^2 = \sinh r \sin \phi$ $(ds^2)_1 = dr^2 + \sinh^2 r d\phi^2$ $\Gamma^r_{\phi\phi} = -\sinh r \cosh r$ $\Gamma^\phi_{\phi r} = 1/\tanh r$ $dS = \sinh r dr \wedge d\phi$
$\mathbf{NH}^{1+1}_+ = S^2_{[+1/\tau^2], 0}$ $x^0 = \cos(r/\tau)$ $x^1 = \tau \sin(r/\tau)$ $x^2 = \tau \sin(r/\tau) \chi$ $(ds^2)_1 = dr^2$ $(ds^2)_2 = \tau^2 \sin^2(r/\tau) d\chi^2 \quad r = r_0$ $\Gamma^r_{\chi\chi} = 0 \quad \Gamma^\chi_{\chi r} = \frac{1}{\tau \tan(r/\tau)}$ $dS = \tau \sin(r/\tau) dr \wedge d\chi$	$\mathbf{G}^{1+1} = S^2_{[0], 0}$ $x^0 = 1$ $x^1 = r$ $x^2 = r \chi$ $(ds^2)_1 = dr^2$ $(ds^2)_2 = r^2 d\chi^2 \quad r = r_0$ $\Gamma^r_{\chi\chi} = 0 \quad \Gamma^\chi_{\chi r} = 1/r$ $dS = r dr \wedge d\chi$	$\mathbf{NH}^{1+1}_- = S^2_{[-1/\tau^2], 0}$ $x^0 = \cosh(r/\tau)$ $x^1 = \tau \sinh(r/\tau)$ $x^2 = \tau \sinh(r/\tau) \chi$ $(ds^2)_1 = dr^2$ $(ds^2)_2 = \tau^2 \sinh^2(r/\tau) d\chi^2 \quad r = r_0$ $\Gamma^r_{\chi\chi} = 0 \quad \Gamma^\chi_{\chi r} = \frac{1}{\tau \tanh(r/\tau)}$ $dS = \tau \sinh(r/\tau) dr \wedge d\chi$
$\mathbf{AdS}^{1+1} = S^2_{[+1/\tau^2], -1/c^2}$ $x^0 = \cos(r/\tau)$ $x^1 = \tau \sin(r/\tau) \cosh(\chi/c)$ $x^2 = c\tau \sin(r/\tau) \sinh(\chi/c)$ $(ds^2)_1 = dr^2 - \frac{\tau^2}{c^2} \sin^2(r/\tau) d\chi^2$ $\Gamma^r_{\chi\chi} = \frac{\tau}{c^2} \sin(r/\tau) \cos(r/\tau)$ $\Gamma^\chi_{\chi r} = \frac{1}{\tau \tan(r/\tau)}$ $dS = \tau \sin(r/\tau) dr \wedge d\chi$	$\mathbf{M}^{1+1} = S^2_{[0], -1/c^2}$ $x^0 = 1$ $x^1 = r \cosh(\chi/c)$ $x^2 = c r \sinh(\chi/c)$ $(ds^2)_1 = dr^2 - \frac{1}{c^2} r^2 d\chi^2$ $\Gamma^r_{\chi\chi} = \frac{1}{c^2} r$ $\Gamma^\chi_{\chi r} = 1/r$ $dS = r dr \wedge d\chi$	$\mathbf{dS}^{1+1} = S^2_{[-1/\tau^2], -1/c^2}$ $x^0 = \cosh(r/\tau)$ $x^1 = \tau \sinh(r/\tau) \cosh(\chi/c)$ $x^2 = c\tau \sinh(r/\tau) \sinh(\chi/c)$ $(ds^2)_1 = dr^2 - \frac{\tau^2}{c^2} \sinh^2(r/\tau) d\chi^2$ $\Gamma^r_{\chi\chi} = \frac{\tau}{c^2} \sinh(r/\tau) \cosh(r/\tau)$ $\Gamma^\chi_{\chi r} = \frac{1}{\tau \tanh(r/\tau)}$ $dS = \tau \sinh(r/\tau) dr \wedge d\chi$

Table 5: Generic form for the equations of cycles, (including as particular cases geodesics, equidistants and circles) in the spaces $S^2_{[\kappa_1], \kappa_2}$ in the three geodesic coordinate systems. The circle equations can also be read as expressions for the finite (time-like) distance between two points, and the equidistant equations as expressions for the finite (time-like) distance between a point and a (space-like) line.

Geodesic parallel I coordinates (a, y)	
Cycles	$\alpha_0 C_{\kappa_1}(a) + \alpha_1 S_{\kappa_1}(a) = \frac{\alpha}{C_{\kappa_1 \kappa_2}(y)} - \alpha_2 T_{\kappa_1 \kappa_2}(y) = \alpha C_{-\kappa_1 \kappa_2}(y^\wedge) - \alpha_2 S_{-\kappa_1 \kappa_2}(y^\wedge)$ $= \left(\frac{\alpha \sqrt{\kappa_1 \kappa_2} - \alpha_2}{2\sqrt{\kappa_1 \kappa_2}} \right) \left(\frac{1 + \sqrt{\kappa_1 \kappa_2} T_{\kappa_1 \kappa_2}(y/2)}{1 - \sqrt{\kappa_1 \kappa_2} T_{\kappa_1 \kappa_2}(y/2)} \right) + \left(\frac{\alpha \sqrt{\kappa_1 \kappa_2} + \alpha_2}{2\sqrt{\kappa_1 \kappa_2}} \right) \left(\frac{1 - \sqrt{\kappa_1 \kappa_2} T_{\kappa_1 \kappa_2}(y/2)}{1 + \sqrt{\kappa_1 \kappa_2} T_{\kappa_1 \kappa_2}(y/2)} \right)$
Geodesics	$T_{\kappa_1 \kappa_2}(y) = \beta_0 C_{\kappa_1}(a) + \beta_1 S_{\kappa_1}(a) \quad \text{and} \quad a = a_0$
Equidistants	$S_{\kappa_1}^2(d) = \kappa_2 \frac{\{S_{\kappa_1 \kappa_2}(y) - C_{\kappa_1 \kappa_2}(y)(\beta_0 C_{\kappa_1}(a) + \beta_1 S_{\kappa_1}(a))\}^2}{1 + \kappa_2 \beta_1^2 + \kappa_1 \kappa_2 \beta_0^2} \quad (\kappa_2 \neq 0)$
Circles	$C_{\kappa_1}(\rho) = C_{\kappa_1 \kappa_2}(y) C_{\kappa_1 \kappa_2}(y_0) C_{\kappa_1}(a - a_0) + \kappa_1 \kappa_2 S_{\kappa_1 \kappa_2}(y) S_{\kappa_1 \kappa_2}(y_0)$ $V_{\kappa_1}(\rho) = C_{\kappa_1 \kappa_2}(y) C_{\kappa_1 \kappa_2}(y_0) V_{\kappa_1}(a - a_0) + \kappa_2 V_{\kappa_1 \kappa_2}(y - y_0)$
Geodesic parallel II coordinates (x, b)	
Cycles	$\alpha_0 C_{\kappa_1 \kappa_2}(b) + \alpha_2 S_{\kappa_1 \kappa_2}(b) = \frac{\alpha}{C_{\kappa_1}(x)} - \alpha_1 T_{\kappa_1}(x) = \alpha C_{-\kappa_1}(x^\wedge) - \alpha_1 S_{-\kappa_1}(x^\wedge)$ $= \left(\frac{\alpha \sqrt{\kappa_1} - \alpha_1}{2\sqrt{\kappa_1}} \right) \left(\frac{1 + \sqrt{\kappa_1} T_{\kappa_1}(x/2)}{1 - \sqrt{\kappa_1} T_{\kappa_1}(x/2)} \right) + \left(\frac{\alpha \sqrt{\kappa_1} + \alpha_1}{2\sqrt{\kappa_1}} \right) \left(\frac{1 - \sqrt{\kappa_1} T_{\kappa_1}(x/2)}{1 + \sqrt{\kappa_1} T_{\kappa_1}(x/2)} \right)$
Geodesics	$T_{\kappa_1}(x) = \beta_0 C_{\kappa_1 \kappa_2}(b) + \beta_2 S_{\kappa_1 \kappa_2}(b) \quad \text{and} \quad b = b_0$
Equidistants	$S_{\kappa_1}^2(d) = \kappa_2 \frac{\{S_{\kappa_1}(x) - C_{\kappa_1}(x)(\beta_0 C_{\kappa_1 \kappa_2}(b) + \beta_2 S_{\kappa_1 \kappa_2}(b))\}^2}{\kappa_2 + \beta_2^2 + \kappa_1 \kappa_2 \beta_0^2} \quad (\kappa_2 \neq 0)$
Circles	$C_{\kappa_1}(\rho) = C_{\kappa_1}(x) C_{\kappa_1}(x_0) C_{\kappa_1 \kappa_2}(b - b_0) + \kappa_1 S_{\kappa_1}(x) S_{\kappa_1}(x_0)$ $V_{\kappa_1}(\rho) = \kappa_2 C_{\kappa_1}(x) C_{\kappa_1}(x_0) V_{\kappa_1 \kappa_2}(b - b_0) + V_{\kappa_1}(x - x_0)$
Geodesic polar coordinates (r, ϕ)	
Cycles	$\alpha_1 C_{\kappa_2}(\phi) + \alpha_2 S_{\kappa_2}(\phi) = \frac{\alpha}{S_{\kappa_1}(r)} - \frac{\alpha_0}{T_{\kappa_1}(r)} = \kappa_1 \left(\frac{\alpha + \alpha_0}{2} \right) T_{\kappa_1}(r/2) + \left(\frac{\alpha - \alpha_0}{2} \right) \frac{1}{T_{\kappa_1}(r/2)}$
Geodesics	$\frac{1}{T_{\kappa_1}(r)} = \beta_1 C_{\kappa_2}(\phi) + \beta_2 S_{\kappa_2}(\phi) \quad \text{and} \quad \phi = \phi_0$
Equidistants	$S_{\kappa_1}^2(d) = \kappa_2 \frac{\{C_{\kappa_1}(r) - S_{\kappa_1}(r)(\beta_1 C_{\kappa_2}(\phi) + \beta_2 S_{\kappa_2}(\phi))\}^2}{\beta_2^2 + \kappa_2 \beta_1^2 + \kappa_1 \kappa_2} \quad (\kappa_2 \neq 0)$
Circles	$C_{\kappa_1}(\rho) = C_{\kappa_1}(r) C_{\kappa_1}(r_0) + \kappa_1 S_{\kappa_1}(r) S_{\kappa_1}(r_0) C_{\kappa_2}(\phi - \phi_0)$ $V_{\kappa_1}(\rho) = V_{\kappa_1}(r - r_0) + \kappa_2 S_{\kappa_1}(r) S_{\kappa_1}(r_0) V_{\kappa_2}(\phi - \phi_0)$

Table 6: Particularized equations for geodesics (besides $a = a_0$) and circles (constant geodesic ‘time-like’ distance ρ to a fixed center (a_0, y_0)) for the nine spaces $S_{[\kappa_1], \kappa_2}^2$ in geodesic parallel I coordinates and with the same conventions as in table 3.

$\mathbf{S}^2 = S_{[+],+}^2$	$\mathbf{E}^2 = S_{[0],+}^2$	$\mathbf{H}^2 = S_{[-],+}^2$
$\tan y = \beta_0 \cos a + \beta_1 \sin a$	$y = \beta_0 + \beta_1 a$	$\tanh y = \beta_0 \cosh a + \beta_1 \sinh a$
$\cos \rho = \cos y \cos y_0 \cos(a - a_0)$ $+ \sin y \sin y_0$	$\rho^2 = (a - a_0)^2$ $+ (y - y_0)^2$	$\cosh \rho = \cosh y \cosh y_0 \cosh(a - a_0)$ $- \sinh y \sinh y_0$
$\mathbf{NH}_+^{1+1} = S_{[+1/\tau^2],0}^2$	$\mathbf{G}^{1+1} = S_{[0],0}^2$	$\mathbf{NH}_-^{1+1} = S_{[-1/\tau^2],0}^2$
$y = \beta_0 \cos(t/\tau) + \beta_1 \tau \sin(t/\tau)$	$y = \beta_0 + \beta_1 t$	$y = \beta_0 \cosh(t/\tau) + \beta_1 \tau \sinh(t/\tau)$
$\cos(\rho/\tau) = \cos((t - t_0)/\tau)$	$\rho^2 = (t - t_0)^2$	$\cosh(\rho/\tau) = \cosh((t - t_0)/\tau)$
$\mathbf{AdS}^{1+1} = S_{[+1/\tau^2],-1/c^2}^2$	$\mathbf{M}^{1+1} = S_{[0],-1/c^2}^2$	$\mathbf{dS}^{1+1} = S_{[-1/\tau^2],-1/c^2}^2$
$c\tau \tanh(y/c\tau) = \beta_0 \cos(t/\tau) + \beta_1 \tau \sin(t/\tau)$	$y = \beta_0 + \beta_1 t$	$c\tau \tan(y/c\tau) = \beta_0 \cosh(t/\tau) + \beta_1 \tau \sinh(t/\tau)$
$\cos(\rho/\tau) = -\sinh(y/c\tau) \sinh(y_0/c\tau)$ $+ \cosh(y/c\tau) \cosh(y_0/c\tau) \cos((t - t_0)/\tau)$	$\rho^2 = (t - t_0)^2$ $- \frac{1}{c^2}(y - y_0)^2$	$\cosh(\rho/\tau) = \sin(y/c\tau) \sin(y_0/c\tau)$ $+ \cos(y/c\tau) \cos(y_0/c\tau) \cosh((t - t_0)/\tau)$

Table 7: Parametrized differential realizations for the generators $\{P_i, J_{12}, D, L_i, G_i, R_i\}$ of conformal algebras $\text{conf}_{\kappa_1, \kappa_2}$. Note the three pairs $P_1, L_2; P_2, L_1; J, D$ with a specially simple expression in each of the three coordinate systems.

Geodesic parallel I coordinates (a, y)	
$P_1 = -\partial_a$	$P_2 = -\kappa_1 \kappa_2 S_{\kappa_1}(a) T_{\kappa_1 \kappa_2}(y) \partial_a - C_{\kappa_1}(a) \partial_y$
$J_{12} = \kappa_2 C_{\kappa_1}(a) T_{\kappa_1 \kappa_2}(y) \partial_a - S_{\kappa_1}(a) \partial_y$	$D = -\frac{S_{\kappa_1}(a)}{C_{\kappa_1 \kappa_2}(y)} \partial_a - C_{\kappa_1}(a) S_{\kappa_1 \kappa_2}(y) \partial_y$
$L_1 = -\frac{C_{\kappa_1}(a)}{C_{\kappa_1 \kappa_2}(y)} \partial_a + \kappa_1 S_{\kappa_1}(a) S_{\kappa_1 \kappa_2}(y) \partial_y$	$L_2 = -C_{\kappa_1 \kappa_2}(y) \partial_y$
$G_1 = \frac{1}{C_{\kappa_1 \kappa_2}(y)} (V_{\kappa_1}(a) - \kappa_2 V_{\kappa_1 \kappa_2}(y)) \partial_a + S_{\kappa_1}(a) S_{\kappa_1 \kappa_2}(y) \partial_y$	
$G_2 = \kappa_2 S_{\kappa_1}(a) T_{\kappa_1 \kappa_2}(y) \partial_a - (V_{\kappa_1}(a) - \kappa_2 V_{\kappa_1 \kappa_2}(y)) \partial_y$	
$R_1 = -\frac{1}{2} \left(1 + \frac{C_{\kappa_1}(a)}{C_{\kappa_1 \kappa_2}(y)} \right) \partial_a + \frac{1}{2} \kappa_1 S_{\kappa_1}(a) S_{\kappa_1 \kappa_2}(y) \partial_y$	
$R_2 = -\frac{1}{2} \kappa_1 \kappa_2 S_{\kappa_1}(a) T_{\kappa_1 \kappa_2}(y) \partial_a - \frac{1}{2} (C_{\kappa_1}(a) + C_{\kappa_1 \kappa_2}(y)) \partial_y$	
Geodesic parallel II coordinates (x, b)	
$P_1 = -C_{\kappa_1 \kappa_2}(b) \partial_x - \kappa_1 T_{\kappa_1}(x) S_{\kappa_1 \kappa_2}(b) \partial_b$	$P_2 = -\partial_b$
$J_{12} = \kappa_2 S_{\kappa_1 \kappa_2}(b) \partial_x - T_{\kappa_1}(x) C_{\kappa_1 \kappa_2}(b) \partial_b$	$D = -S_{\kappa_1}(x) C_{\kappa_1 \kappa_2}(b) \partial_x - \frac{S_{\kappa_1 \kappa_2}(b)}{C_{\kappa_1}(x)} \partial_b$
$L_1 = -C_{\kappa_1}(x) \partial_x$	$L_2 = \kappa_1 \kappa_2 S_{\kappa_1}(x) S_{\kappa_1 \kappa_2}(b) \partial_x - \frac{C_{\kappa_1 \kappa_2}(b)}{C_{\kappa_1}(x)} \partial_b$
$G_1 = (V_{\kappa_1}(x) - \kappa_2 V_{\kappa_1 \kappa_2}(b)) \partial_x + T_{\kappa_1}(x) S_{\kappa_1 \kappa_2}(b) \partial_b$	
$G_2 = \kappa_2 S_{\kappa_1}(x) S_{\kappa_1 \kappa_2}(b) \partial_x - \frac{1}{C_{\kappa_1}(x)} (V_{\kappa_1}(x) - \kappa_2 V_{\kappa_1 \kappa_2}(b)) \partial_b$	
$R_1 = -\frac{1}{2} (C_{\kappa_1}(x) + C_{\kappa_1 \kappa_2}(b)) \partial_x - \frac{1}{2} \kappa_1 T_{\kappa_1}(x) S_{\kappa_1 \kappa_2}(b) \partial_b$	
$R_2 = \frac{1}{2} \kappa_1 \kappa_2 S_{\kappa_1}(x) S_{\kappa_1 \kappa_2}(b) \partial_x - \frac{1}{2} \left(1 + \frac{C_{\kappa_1 \kappa_2}(b)}{C_{\kappa_1}(x)} \right) \partial_b$	
Geodesic polar coordinates (r, ϕ)	
$P_1 = -C_{\kappa_2}(\phi) \partial_r + \frac{S_{\kappa_2}(\phi)}{T_{\kappa_1}(r)} \partial_\phi$	$P_2 = -\kappa_2 S_{\kappa_2}(\phi) \partial_r - \frac{C_{\kappa_2}(\phi)}{T_{\kappa_1}(r)} \partial_\phi$
$J_{12} = -\partial_\phi$	$D = -S_{\kappa_1}(r) \partial_r$
$L_1 = -C_{\kappa_2}(\phi) C_{\kappa_1}(r) \partial_r + \frac{S_{\kappa_2}(\phi)}{S_{\kappa_1}(r)} \partial_\phi$	$L_2 = -\kappa_2 S_{\kappa_2}(\phi) C_{\kappa_1}(r) \partial_r - \frac{C_{\kappa_2}(\phi)}{S_{\kappa_1}(r)} \partial_\phi$
$G_1 = C_{\kappa_2}(\phi) V_{\kappa_1}(r) \partial_r + S_{\kappa_2}(\phi) \frac{V_{\kappa_1}(r)}{S_{\kappa_1}(r)} \partial_\phi$	
$G_2 = \kappa_2 S_{\kappa_2}(\phi) V_{\kappa_1}(r) \partial_r - C_{\kappa_2}(\phi) \frac{V_{\kappa_1}(r)}{S_{\kappa_1}(r)} \partial_\phi$	
$R_1 = -\frac{1}{2 T_{\kappa_1}(r/2)} (C_{\kappa_2}(\phi) S_{\kappa_1}(r) \partial_r - S_{\kappa_2}(\phi) \partial_\phi)$	
$R_2 = -\frac{1}{2 T_{\kappa_1}(r/2)} (\kappa_2 S_{\kappa_2}(\phi) S_{\kappa_1}(r) \partial_r + C_{\kappa_2}(\phi) \partial_\phi)$	

Table 8: The Laplace–Beltrami operator $\mathcal{C}$ giving rise to differential Laplace and wave-type equations $\mathcal{C}\Phi = 0$ in geodesic parallel I $\Phi(a, y) \equiv \Phi(t, y)$ and polar $\Phi(r, \phi) \equiv \Phi(r, \chi)$ coordinates for the nine CK spaces. This operator has the corresponding conformal algebra symmetry, $\text{conf}_{\kappa_1, \kappa_2}$, which is independent of κ_1 and isomorphic to either $so(3, 1)$, $iso(2, 1)$ or $so(2, 2)$; this algebra appears as a heading in each entry.

$so(3, 1) : \mathbf{S}^2 = S_{[+],+}^2$ $\frac{1}{\cos^2 y} \partial_a^2 + \partial_y^2 - \tanh y \partial_y$ $\frac{1}{\sin^2 r} \partial_\phi^2 + \partial_r^2 + \frac{1}{\tan r} \partial_r$	$so(3, 1) : \mathbf{E}^2 = S_{[0],+}^2$ $\partial_a^2 + \partial_y^2$ $\frac{1}{r^2} \partial_\phi^2 + \partial_r^2 + \frac{1}{r} \partial_r$	$so(3, 1) : \mathbf{H}^2 = S_{[-],+}^2$ $\frac{1}{\cosh^2 y} \partial_a^2 + \partial_y^2 + \tanh y \partial_y$ $\frac{1}{\sinh^2 r} \partial_\phi^2 + \partial_r^2 + \frac{1}{\tanh r} \partial_r$
$iso(2, 1) : \mathbf{NH}_+^{1+1} = S_{[+1/\tau^2],0}^2$ ∂_y^2 $\frac{1}{\tau^2 \sin^2(r/\tau)} \partial_\chi^2$	$iso(2, 1) : \mathbf{G}^{1+1} = S_{[0],0}^2$ ∂_y^2 $\frac{1}{r^2} \partial_\chi^2$	$iso(2, 1) : \mathbf{NH}_-^{1+1} = S_{[-1/\tau^2],0}^2$ ∂_y^2 $\frac{1}{\tau^2 \sinh^2(r/\tau)} \partial_\chi^2$
$so(2, 2) : \mathbf{AdS}^{1+1} = S_{[+1/\tau^2],-1/c^2}^2$ $\frac{-1}{c^2 \cosh^2(y/c\tau)} \partial_t^2 + \partial_y^2 + \frac{\tanh(y/c\tau)}{c\tau} \partial_y$ $\frac{1}{\tau^2 \sin^2(r/\tau)} \partial_\chi^2 - \frac{1}{c^2} \partial_r^2 - \frac{1}{c^2 \tau \tanh(r/\tau)} \partial_r$	$so(2, 2) : \mathbf{M}^{1+1} = S_{[0],-1/c^2}^2$ $-\frac{1}{c^2} \partial_t^2 + \partial_y^2$ $\frac{1}{r^2} \partial_\chi^2 - \frac{1}{c^2} \partial_r^2 - \frac{1}{c^2 r} \partial_r$	$so(2, 2) : \mathbf{dS}^{1+1} = S_{[-1/\tau^2],-1/c^2}^2$ $\frac{-1}{c^2 \cos^2(y/c\tau)} \partial_t^2 + \partial_y^2 - \frac{\tanh(y/c\tau)}{c\tau} \partial_y$ $\frac{1}{\tau^2 \sinh^2(r/\tau)} \partial_\chi^2 - \frac{1}{c^2} \partial_r^2 - \frac{1}{c^2 \tau \tanh(r/\tau)} \partial_r$

Table 9: Parametrized expressions for the conformal embedding $S_{[\kappa_1],\kappa_2}^2 \mapsto C_{[\kappa_1],\kappa_2}^2$ in three coordinate systems. Coordinates (s^+, s^-, s^1, s^2) parametrize the conformal ambient space, and hence the cone Γ_0 , while $(\tilde{s}^+, \tilde{s}^- = -1, \tilde{s}^1, \tilde{s}^2)$ are a parametrization of the conformal space itself $C_{[\kappa_1],\kappa_2}^2 \equiv \tilde{S}_{[+],\kappa_2}^2$. We have introduced the shorthand notation $\eta_+ = \frac{1+\kappa_1\ell^2}{2\kappa_1\ell^2}$ and $\eta_- = \frac{1-\kappa_1\ell^2}{2\kappa_1\ell^2}$.

Parallel I coordinates	Parallel II coordinates	Polar coordinates
$(a, y) \in S_{[\kappa_1],\kappa_2}^2 \rightarrow (A, Y) \in C_{[\kappa_1],\kappa_2}^2$	$(x, b) \in S_{[\kappa_1],\kappa_2}^2 \rightarrow (X, B) \in C_{[\kappa_1],\kappa_2}^2$	$(r, \phi) \in S_{[\kappa_1],\kappa_2}^2 \rightarrow (R, \Phi) \in C_{[\kappa_1],\kappa_2}^2$
$s^\pm = \pm 1 - \eta_\pm \{1 - C_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y)\}$ $s^1 = S_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y)/\ell$ $s^2 = S_{\kappa_1\kappa_2}(y)/\ell$	$s^\pm = \pm 1 - \eta_\pm \{1 - C_{\kappa_1}(x)C_{\kappa_1\kappa_2}(b)\}$ $s^1 = S_{\kappa_1}(x)/\ell$ $s^2 = C_{\kappa_1}(x)S_{\kappa_1\kappa_2}(b)/\ell$	$s^\pm = \pm 1 - \eta_\pm (1 - C_{\kappa_1}(r))$ $s^1 = S_{\kappa_1}(r)C_{\kappa_2}(\phi)/\ell$ $s^2 = S_{\kappa_1}(r)S_{\kappa_2}(\phi)/\ell$
$\tilde{s}^+ = \frac{1 - \eta_+ \{1 - C_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y)\}}{1 + \eta_- \{1 - C_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y)\}}$ $\tilde{s}^1 = \frac{S_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y)/\ell}{1 + \eta_- \{1 - C_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y)\}}$ $\tilde{s}^2 = \frac{S_{\kappa_1\kappa_2}(y)/\ell}{1 + \eta_- \{1 - C_{\kappa_1}(a)C_{\kappa_1\kappa_2}(y)\}}$	$\tilde{s}^+ = \frac{1 - \eta_+ \{1 - C_{\kappa_1}(x)C_{\kappa_1\kappa_2}(b)\}}{1 + \eta_- \{1 - C_{\kappa_1}(x)C_{\kappa_1\kappa_2}(b)\}}$ $\tilde{s}^1 = \frac{S_{\kappa_1}(x)/\ell}{1 + \eta_- \{1 - C_{\kappa_1}(x)C_{\kappa_1\kappa_2}(b)\}}$ $\tilde{s}^2 = \frac{C_{\kappa_1}(x)S_{\kappa_1\kappa_2}(b)/\ell}{1 + \eta_- \{1 - C_{\kappa_1}(x)C_{\kappa_1\kappa_2}(b)\}}$	$\tilde{s}^+ = \frac{\ell^2 - T_{\kappa_1}^2(r/2)}{\ell^2 + T_{\kappa_1}^2(r/2)}$ $\tilde{s}^1 = \frac{2\ell T_{\kappa_1}(r/2)C_{\kappa_2}(\phi)}{\ell^2 + T_{\kappa_1}^2(r/2)}$ $\tilde{s}^2 = \frac{2\ell T_{\kappa_1}(r/2)S_{\kappa_2}(\phi)}{\ell^2 + T_{\kappa_1}^2(r/2)}$
$\tan A = \frac{T_{\kappa_1}(a)}{\ell} \frac{1 - \kappa_1 T_{\kappa_1}^2(r/2)}{1 - \frac{1}{\ell^2} T_{\kappa_1}^2(r/2)}$ $S_{\kappa_2}(Y) = \frac{S_{\kappa_1\kappa_2}(y)}{\ell} \frac{1 + \kappa_1 T_{\kappa_1}^2(r/2)}{1 + \frac{1}{\ell^2} T_{\kappa_1}^2(r/2)}$	$\sin X = \frac{S_{\kappa_1}(x)}{\ell} \frac{1 + \kappa_1 T_{\kappa_1}^2(r/2)}{1 + \frac{1}{\ell^2} T_{\kappa_1}^2(r/2)}$ $T_{\kappa_2}(B) = \frac{T_{\kappa_1\kappa_2}(b)}{\ell} \frac{1 - \kappa_1 T_{\kappa_1}^2(r/2)}{1 - \frac{1}{\ell^2} T_{\kappa_1}^2(r/2)}$	$\tan^2(R/2) = \frac{1}{\ell^2} T_{\kappa_1}^2(r/2)$ $\Phi = \phi$

Table 10: The conformal embedding in parallel I coordinates for the nine CK spaces and for their compactifications: $(a, y) \equiv (t, y) \in S_{[\kappa_1], \kappa_2}^2 \mapsto (A, Y) \equiv (T, Y) \in C_{[\kappa_1], \kappa_2}^2 \equiv \widetilde{S}_{[+], \kappa_2}^2$. The length ℓ is chosen in such a manner that $\kappa_1 \ell^2 \in \{1, 0, -1\}$ and $\kappa_2 \in \{1, 0, -1/c^2\}$.

$\mathbf{S}^2 \mapsto \mathbf{S}^2$ $\kappa_1 \ell^2 = 1, \kappa_2 = 1$	$\mathbf{E}^2 \mapsto \mathbf{S}^2$ $\kappa_1 \ell^2 = 0, \kappa_2 = 1$	$\mathbf{H}^2 \mapsto \mathbf{S}^2$ $\kappa_1 \ell^2 = -1, \kappa_2 = 1$
$\tilde{s}^+ = \cos(a/\ell) \cos(y/\ell)$	$\tilde{s}^+ = \frac{4 - \{(a/\ell)^2 + (y/\ell)^2\}}{4 + \{(a/\ell)^2 + (y/\ell)^2\}}$	$\tilde{s}^+ = \frac{1}{\cosh(a/\ell) \cosh(y/\ell)}$
$\tilde{s}^1 = \sin(a/\ell) \cos(y/\ell)$	$\tilde{s}^1 = \frac{4a/\ell}{4 + \{(a/\ell)^2 + (y/\ell)^2\}}$	$\tilde{s}^1 = \tanh(a/\ell)$
$\tilde{s}^2 = \sin(y/\ell)$	$\tilde{s}^2 = \frac{4y/\ell}{4 + \{(a/\ell)^2 + (y/\ell)^2\}}$	$\tilde{s}^2 = \frac{\tanh(y/\ell)}{\cosh(a/\ell)}$
$\tan A = \tan(a/\ell)$	$\tan A = \frac{4a/\ell}{4 - \{(a/\ell)^2 + (y/\ell)^2\}}$	$\tan A = \sinh(a/\ell) \cosh(y/\ell)$
$\sin Y = \sin(y/\ell)$	$\sin Y = \frac{4y/\ell}{4 + \{(a/\ell)^2 + (y/\ell)^2\}}$	$\sin Y = \frac{\tanh(y/\ell)}{\cosh(a/\ell)}$
$\mathbf{NH}_+^{1+1} \mapsto \widetilde{\mathbf{NH}}_+^{1+1}$ $\kappa_1 \ell^2 = 1, \kappa_2 = 0$	$\mathbf{G}^{1+1} \mapsto \widetilde{\mathbf{NH}}_+^{1+1}$ $\kappa_1 \ell^2 = 0, \kappa_2 = 0$	$\mathbf{NH}_-^{1+1} \mapsto \widetilde{\mathbf{NH}}_+^{1+1}$ $\kappa_1 \ell^2 = -1, \kappa_2 = 0$
$\tilde{s}^+ = \cos(t/\ell)$	$\tilde{s}^+ = \frac{4 - (t/\ell)^2}{4 + (t/\ell)^2}$	$\tilde{s}^+ = \frac{1}{\cosh(t/\ell)}$
$\tilde{s}^1 = \sin(t/\ell)$	$\tilde{s}^1 = \frac{4t/\ell}{4 + (t/\ell)^2}$	$\tilde{s}^1 = \tanh(t/\ell)$
$\tilde{s}^2 = y/\ell$	$\tilde{s}^2 = \frac{4y/\ell}{4 + (t/\ell)^2}$	$\tilde{s}^2 = \frac{y/\ell}{\cosh(t/\ell)}$
$\tan T = \tan(t/\ell)$	$\tan T = \frac{4t/\ell}{4 - (t/\ell)^2}$	$\tan T = \sinh(t/\ell)$
$Y = y/\ell$	$Y = \frac{4y/\ell}{4 + (t/\ell)^2}$	$Y = \frac{y/\ell}{\cosh(t/\ell)}$
$\mathbf{AdS}^{1+1} \mapsto \widetilde{\mathbf{AdS}}^{1+1}$ $\kappa_1 \ell^2 = 1, \kappa_2 = -1/c^2$	$\mathbf{M}^{1+1} \mapsto \widetilde{\mathbf{AdS}}^{1+1}$ $\kappa_1 \ell^2 = 0, \kappa_2 = -1/c^2$	$\mathbf{dS}^{1+1} \mapsto \widetilde{\mathbf{AdS}}^{1+1}$ $\kappa_1 \ell^2 = -1, \kappa_2 = -1/c^2$
$\tilde{s}^+ = \cos(t/\ell) \cosh(y/c\ell)$	$\tilde{s}^+ = \frac{4 - \{(t/\ell)^2 - (y/c\ell)^2\}}{4 + \{(t/\ell)^2 - (y/c\ell)^2\}}$	$\tilde{s}^+ = \frac{1}{\cosh(t/\ell) \cos(y/c\ell)}$
$\tilde{s}^1 = \sin(t/\ell) \cosh(y/c\ell)$	$\tilde{s}^1 = \frac{4t/\ell}{4 + \{(t/\ell)^2 - (y/c\ell)^2\}}$	$\tilde{s}^1 = \tanh(t/\ell)$
$\tilde{s}^2 = c \sinh(y/c\ell)$	$\tilde{s}^2 = \frac{4y/\ell}{4 + \{(t/\ell)^2 - (y/c\ell)^2\}}$	$\tilde{s}^2 = \frac{c \tanh(y/c\ell)}{\cosh(t/\ell)}$
$\tan T = \tan(t/\ell)$	$\tan T = \frac{4t/\ell}{4 - \{(t/\ell)^2 - (y/c\ell)^2\}}$	$\tan T = \sinh(t/\ell) \cos(y/c\ell)$
$\sinh(Y/c) = \sinh(y/c\ell)$	$\sinh(Y/c) = \frac{4(y/c\ell)}{4 + \{(t/\ell)^2 - (y/c\ell)^2\}}$	$\sinh(Y/c) = \frac{\tanh(y/c\ell)}{\cosh(t/\ell)}$